

Научном већу Института за физику у Београду

Београд, 19.2.2024.

Предмет:

Молба за покретање поступка за стицање звања истраживач-сарадник

На основу испуњености услова предвиђених Правилником о стицању истраживачких и научних звања, прописаног од стране ресорног Министарства, молим Научно веће института за физику у Београду да покрене поступак за мој избор у звање истраживач сарадник.

У прилогу достављам:

1. Мишљење руководиоца лабораторије са предлогом комисије за избор у звање
2. Стручну биографију
3. Преглед научних активности
4. Списак научних радова
5. Потврду о статусу студента докторских студија и пријављеној теми докторске дисертације
6. Копије диплома основних и мастер студија
7. Копије научних радова

С поштовањем,
Ива Росић
истраживач приправник



Научном већу Института за физику у Београду
Београд, 19.2.2024.


Предмет: Мишљење руководиоца лабораторије о избору Иве Росић у звање истраживач сарадник

Ива Росић је запослена на Институту за физику у Београду. Израду докторске дисертације обавља на Катедри за микробиологију Биолошког факултета Универзитета у Београду под руководством др Оље Медић. С обзиром да испуњава све предвиђене услове у складу са Законом о науци и истраживањима и Правилником о стицању истраживачких и научних звања Министарства науке, технолошког развоја и иновација, сагласан сам са покретањем поступка за избор Иве Росић у звање истраживач сарадник.

За састав комисије за избор Иве Росић у звање истраживач сарадник предлажем:

- (1) др Тијана Томашевић-Илић, научни сарадник, Институт за физику у Београду
- (2) др Наташа Томић, научни сарадник, Институт за физику у Београду
- (3) др Тања Берић, редовни професор Биолошког факултета Универзитета у Београду




др Александар Богојевић
Научни саветник

Директор Института за физику у Београду

Биографски подаци кандидаткиње

Ива М. Росић рођена је 9.5.1997. у Ужицу. У Ужицу је завршила основну школу као носилац дипломе „Вук Караџић“ и средњу медицинску школу, смер фармацеутски техничар, као ђак генерације. Биолошки факултет Универзитета у Београду, модул Молекуларна биологија и физиологија уписала је школске 2016/17. године, а дипломирала 2020. године са просечном оценом 9,60. Школске 2020/21. уписала је мастер студије на Биолошком факултету Универзитета у Београду, модул Биологија микроорганизама, које је завршила са просечном оценом 10,00. Мастер рад под насловом: „Генотипизација и одређивање ДНК фрагмената специфичних за сој у колекцији биоконтролних *Bacillus* spp. изолата“ урадила је на Катедри за микробиологију Биолошког факултета под менторством проф. др Славише Станковића и др Ивана Николића. Школске 2021/22. уписала је докторске академске студије на модулу Биологија микроорганизама на Биолошком факултету Универзитета у Београду. Од априла 2022. до маја 2023. била је запослена као истраживач приправник на Катедри за микробиологију Биолошког факултета, Универзитета у Београду, где је учествовала у настави на практичном делу курса из Микробиологије на основним академским студијама. Такође, у школској 2021/22. учествовала је као сарадник у реализацији семинара из Биологије у Истраживачкој станици Петница.

Од јуна 2023. запослена је као истраживач приправник на Институту за физику, Универзитета у Београду. Тренутно учествује на два билатерална пројекта. У оквиру билатералног пројекта са Француском, под називом: „Очување биљног здравља: развој капацитета за теренски надзор *Clavibacter michiganensis* subs. *michiganensis* на усевима парадајза“, била је у једнонедељној истраживачкој посети Националном институту за пољопривреду и животну средину у Авињону (ИНРАЕ), током јуна 2023. године. У оквиру билатералног пројекта са Словенијом, „Стварање капацитета за надзор биљних патогена: Развијање qPCR тестова за детекцију *P. syringae* pv. *aptata* патогена шећерне репе“, била је на једномесечном истраживачком боравку на Националном институту за биологију (НИБ) Љубљани, током новембра 2023. године.

Током јула и августа 2018. године Ива Росић је остварила истраживачки боравак на Институту Гулбенкиан у Лисабону у оквиру летње школе Undergraduate Summer School IGC (Instituto Gulbenkian de Ciencia), а септембра исте године је у организацији Центра за научно-истраживачки рад студената Биолошког факултета радила студентску летњу праксу у Клиничком центру Србије, у Служби за патохистологију. Учествовала је као волонтер у више пројеката, конгреса и организација. Могу се издвојити учешће на курсу Биохемијског друштва Србије: Биохемија у служби здравља- изучавање биохемијских и сродних метода које нуде решења у кризним ситуацијама (мај 2022.), учешће у организацији и реализацији програма за Европску ноћ истраживача 2019. године и чланство у омладинској организацији за заштиту животне средине ЕсоHub као Еко-ментор ученицима основних школа. Ива Росић је, такође била носилац стипендије Фонда за младе таленте- Доситеја за 2019/20. и 2020/21. годину, стипендије града Ужица за талентоване ученике и студенте од 2016. до 2020. године и студентске стипендије Министарства просвете, науке и технолошког развоја.

Преглед научне активности кандидаткиње

У досадашњем раду Ива Росић се бавила испитавањем фенотипских и генотипских карактеристика *Bacillus* spp. изолата значајних у колонизацији биљних ткива, биоконтролној активности и особинама значајним у поспешивању раста баљака. Способност биоконтролне активности праћена је против биљног патогена *Pseudomonas syringae* pv. *aptata*, изазовача лисне пегавости шећерне репе.

Бактерије из рода *Bacillus* су међу највише истраживаним и најчешће примењиваним агенсима за биоконтрол и поспешивање раста биљака. Овај род обухвата широко распрострањене, аеробне и факултативно анаеробне, штапићасте, грам-позитивне бактерије са способношћу продукције каталазе. Имају способност производње великог броја ензима који им омогућавају коришћење различитих супстрата за раст. Једна од најзначајнијих особина бактерија из рода *Bacillus* је способност формирања ендоспора. У природи им ова способност омогућава преживљавање неповољних услова животне средине, што им пружа значајну компетитивну предност. Захваљујући ниским нутритивним захтевима, њихово гајење у лабораторијским условима и индустријским биореакторима је једноставно и економично. Формирање ендоспора је итекако значајно у формулацији биоконтролних и препарата за поспешивање раста биљака због стабилности и могућности дуготрајног складиштења. С обзиром на то да су бактерије из рода *Bacillus* доминантан род у ризосфери и филосфери биљака, сматрају се безбедним по живи свет и околину, и погодним за примену у пољопривреди са циљем заштите усева.

Највећи изазов у примени изолата *Bacillus* spp. у биоконтроли и поспешивању раста биљака представља прелаз из контролисаног лабораторијског тестирања на испитивања у пољима. Такође, европске регулативе за комерцијализацију биолошких препарата на бази микроорганизама налажу испитивање постојаности производа у животној средини. Стога је неопходно развити поуздане методе идентификације, квантификације и праћења динамике популације примењених микроорганизама *in situ*. Кандидаткиња у својим истраживањима прати динамику популације биоконтролног соја *in planta* у одсуству и присуству високо-вирулентног фитопатогеног соја *Pseudomonas syringae* pv. *aptata*. Детекција и квантификација одабраних биоконтролних сојева у филосфери шећерне репе омогућена је развојем и амплификацијом сој-специфичних молекуларних маркера (*Sequence Characterized Amplified Region - SCAR*). Упоредо, врши се и процена биоконтролна ефикасност *Bacillus* spp. сојева примењених на два начина, пре и након инокулације патогена (превентивни и куративни ефекат).

Такође, кандидаткиња се бави и израдом специфичних молекуларних метода (*Loop-Mediated Isothermal Amplification - LAMP*) за детекцију карантинског биљног патогена парадајза, *Clavibacter michiganensis* subsp. *michiganensis*. Као и израдом прајмера за специфичну детекцију *P. syringae* сојева из филогенетске групе два, у којој је груписан највећи број изазивача болести код различитих биљних врста.

Списак објављених радова кандидаткиње

Рад у врхинском међународном часопису

1. Anteljević, M., Rosić, I., Medić, O., Kolarević, S., Berić, T., Stanković, S., Nikolić, I. (2023). Occurrence of plant pathogenic *Pseudomonas syringae* in the Danube River Basin: abundance and diversity assessment. *Phytopathology Research*, 5(1). <https://doi.org/10.1186/s42483-023-00174-0>. [M21]

Рад у међународном часопису

1. Rosić, I., Nikolić, I., Ranković, T., Anteljević, M., Medić, O., Berić, T., Stanković, S. (2023). Genotyping-driven diversity assessment of biocontrol potent *Bacillus* spp. Strain collection as a potential method for the development of strain-specific biomarkers. *Archives of Microbiology*, 205(4). <https://doi.org/10.1007/s00203-023-03460-9>. [M23]

Конгресна саопштења са међународног скупа штампана у изводу

1. Rosić, I., Nikolić, I., Ranković, T., Anteljević, M., Berić, T., Stanković, S., Medić, O. (2023, Sep 13-15). Screening of AHL lactonase activity in *Bacillus* spp. strains isolated from different natural samples. ICGEB Workshop Trends in microbial solutions for sustainable agriculture, Belgrade, Serbia, Book of Abstracts, 68. [M34]

2. Anteljević, M., Rosić, I., Ranković, T., Medić, O., Berić, T., Stanković, S., Nikolić, I. (2023, Sep 13-15). The presence of ice nucleation active *Pseudomonas syringae* in the Danube River Basin. ICGEB Workshop Trends in microbial solutions for sustainable agriculture, Belgrade, Serbia, Book of Abstracts, 67. [M34]

3. Rosić, I., Anteljević, M., Ranković, T., Nikolić, I., Stanković, S., Berić, T., Medić, O. (2023, May 15-18). Population dynamics of *Bacillus amyloliquefaciens* SS-38.4 in the phyllosphere of sugar beet and its biocontrol activity against *Pseudomonas syringae* pv. *aptata* P21. 2nd International Molecular Plant Protection Congress, Orhangazi, Turkey, Program and Abstract Book, 113. [M34]

4. Anteljević, M., Rosić, I., Ranković, T., Medić, O., Berić, T., Stanković, S., Nikolić, I. (2023, May 15-18). Phylogenetic analysis of *Pseudomonas syringae* isolates from the Danube River Basin revealed association with past epidemics in Serbia. 2nd International Molecular Plant Protection Congress, Orhangazi, Turkey, Program and Abstract Book, 113. [M34]

5. Nikolić, I., Pavlović, T., **Rosić, I.**, Anteljević, M., Medić, O., Berić, T., Stanković, S. (2022, July 3-8). Phylogenomic status of two *P. syringae* strains P16 and P21 with different pathogenicity isolated from sugar beet in Serbia. 14th International Conference on Plant Pathogenic Bacteria (ICPPB), Assisi, Italy, Book of Abstracts, 129. [M34]

Конгресна саопштења са националног скупа штампана у изводу

1. **Rosić, I.**, Nikolić, I., Medić, O., Pavlović, T., Anteljević, M., Berić, T., Stanković, S. (2022, Sep 21-25). *In vitro* ispitivanje potencijala lipopeptidnih ekstrakata izolata *Bacillus* spp. za suzbijanje biljnog patogena *Pseudomonas syringae*. Treći kongres biologa Srbije, Knjiga sažetaka, 260. [M64]


2. Anteljević, M., Nikolić, I., Kolarević, S., **Rosić, I.**, Pavlović, T., Berić, T., Stanković, S. (2022, Sep 21-25). Zastupljenost biljnog patogena *Pseudomonas syringae* u Uvačkim jezerima. Treći kongres biologa Srbije, Zlatibor, Srbija, Knjiga sažetaka, 263. [M64]

RESEARCH

Open Access



Occurrence of plant pathogenic *Pseudomonas syringae* in the Danube River Basin: abundance and diversity assessment

Marina Anteljević¹, Iva Rosić¹, Olja Medić¹, Stoimir Kolarević², Tanja Berić¹, Slaviša Stanković¹ and Ivan Nikolić^{1*} 

Abstract

Plant pathogenic strains of *Pseudomonas syringae* (*Psy*) spp. have been detected in nonagricultural habitats, including those associated with the water cycle. Their presence in aquatic systems allows dissemination over long distances, especially with irrigation practices. In May 2021, we sampled 15 sites along the Danube River Basin in Serbia to gain insight into *P. syringae* abundance and diversity. We identified 79 *Psy* strains using *Psy*-specific primers, and a partial sequence of the citrate synthase (*cts*) house-keeping gene has served for phylogenetic diversity assessments. Phenotypic diversity determination included characterizing features linked with survival and pathogenic lifestyle. The ice nucleation activity, pectinolytic activity, swimming and swarming assays, and hypersensitive reaction on plants were tested. *Psy* was detected at ten of 15 sites examined at abundance ranging from 1.0×10^2 to 1.2×10^4 CFU/L. We discovered the presence of four phylogroups, with phylogroup 2 being the most abundant, followed by phylogroups 7, 9, and 13. The hypersensitive reaction was induced by 68.63% of the isolates from the collection. A partial sequence comparison of the *cts* gene showed 100% similarity between isolates from cherry plants epidemics in Serbia caused by *Psy* and isolates from the Danube River. Our results suggest that the Danube River, extensively used for irrigation of agricultural fields, harbors diverse strains of *Psy*, which possess various features that could lead to potential disease outbreaks on crops. This study represents the first in-depth analysis of *Psy* abundance and diversity in the Danube River Basin. It sets the ground for future pre-epidemic studies and seasonal monitoring of *Psy* population dynamics.

Keywords *Pseudomonas syringae*, Abundance, Diversity, Phylogeny, Ice nucleation activity, Motility

Background

Pseudomonas syringae (*Psy*) is a well-studied, widespread complex of plant pathogenic bacteria, a causative agent of diseases on many species, including herbaceous and woody plants (Gutiérrez-Barranquero et al. 2019). Symptoms of the diseases caused by this complex of bacteria can vary from mild to severe. Plants can develop localized manifestations, such as leaf spots, blight, specks, and wilting. Some pathogen species can move through vascular tissue and induce systemic disease (Donati et al. 2020). Similar to many other plant pathogens, the evolution of its virulence has been widely investigated from an agro-centric perspective based on interaction

*Correspondence:

Ivan Nikolić
ivan.nikolic@bio.bg.ac.rs

¹ Faculty of Biology, Center for Biological Control and Plant Growth Promotion, University of Belgrade, Studentski Trg 16, Belgrade 11000, Serbia

² Institute for Biological Research 'Siniša Stanković', National Institute of Republic of Serbia, Department of Hydroecology and Water Protection, University of Belgrade, Bulevar Despota Stefana 142, Belgrade 11000, Serbia



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

with its primary hosts (Morris et al. 2009; Scortichini et al. 2012). However, over the past decade, reports have been increasing on the importance of alternative plant hosts and habitats beyond agroecosystems (Morris et al. 2008, 2010). The presence of *P. syringae* was confirmed in habitats correlating to the water cycle, such as rivers, lakes, rainwater, snow, and clouds (Morris et al. 2008). The production of ice-nucleating proteins in some *P. syringae* strains enables them to catalyze ice formation at warmer temperatures than temperatures at which pure water freezes (Maki et al. 1974). It was hypothesized that by catalyzing ice formation in clouds, bacteria provide a deposition to nutrient-rich niches through precipitation (De Araujo et al. 2019). Moreover, ice nucleation activity also threatens plant health, increasing the risk of frost injury (Karimi et al. 2020).

Many of the *P. syringae* hosts are economically important crops, so it is essential to investigate all potential reservoirs of the bacteria. In that context, freshwater habitats are important as they are used for irrigation purposes in agricultural areas and are possible sources of pathogen inoculum. There are many reports about plant pathogenic bacteria in open irrigation systems, such as those from genera *Xanthomonas*, *Pectobacterium*, and *Dickeya* (Faye et al. 2018; Fayette et al. 2018; Hugouvieux-Cotte-Pattat et al. 2019). *P. syringae* was previously detected in freshwater by Morris et al. (2008, 2010), in a freshwater lake in Virginia by Pietsch et al. (2017), also in the study of *Pseudomonas* species diversity along the Danube (Mulet et al. 2020). After cantaloupe blight epidemics caused by *P. syringae* appeared in Southern France in 1993, a post-epidemic study of *P. syringae* in water retention basins was conducted, and the causative agent was isolated from there (Riffaud and Morris 2002). Similarly, for the plant pathogenic fungus *Verticillium dahlia*, irrigation was the primary dispersal means to cause the contamination (Baroudy et al. 2018). Repeated irrigation from the same contaminated water source is an additional danger because the pathogen is reintroduced into the field (Parke et al. 2019). One of the biggest concerns is the emergence of new and highly aggressive populations of plant pathogenic bacteria, considering the enormous genetic diversity in environmental reservoirs. A much broader diversity of plant pathogenic *P. syringae* was obtained from alpine headwaters compared to those isolated from infected plants (Morris et al. 2010; Berge et al. 2014). *P. syringae* environmental isolates, closely related to *P. syringae* pv. *tomato*, had a broader host range but caused almost the same disease symptoms when inoculated in the laboratory under optimal conditions (Monteil et al. 2013). Moreover, *D. dianthicola* isolates from the rivers were

more aggressive on potatoes than strains isolated from diseased plants (Laurila et al. 2008). Environmental strains surrounded by various microbial species have higher chances of acquiring new loci by horizontal gene transfer (Dillon et al. 2019). Insights into the abundance of *P. syringae* in aquatic habitats and assessment of its diversity are important to evaluate its suitability for irrigation. Water inhabited with pathogens is a possible infection source and can lead to epidemics with consequential yield losses (Monteil et al. 2016).

The Danube is the second-longest river in Europe that links more countries than any other river in the world. It flows through or along the borders of ten European countries (Dávid and Madudová 2019). It is the longest river in the Black Sea Basin of Serbia, collecting waters from 92% of Serbian territory. The development of a strategy for irrigation and drainage in the Republic of Serbia is in progress, which will inevitably include the Danube-Tisa-Danube canal network from the territory of Vojvodina. Evaluation of irrigation water quality based on chemical parameters was done for groundwater sources of the Danube (Kurilić et al. 2019) and surface water in the Danube-Tisa-Danube hydrosystem area, encompassing 75% of arable land (Zemunac et al. 2021). Both studies confirmed suitability for irrigation purposes but were based only on chemical characteristics.

Regarding microbial contamination which can potentially cause a severe threat to human health, fecal microbial pollution is a major problem in the Danube River Basin and is the focus of many investigations (Kirschner et al. 2017; Frick et al. 2020; Banciu et al. 2021). If one considers it a potential irrigation source, it is also essential to monitor plant pathogens. To our knowledge, studies of *P. syringae* in the Danube have not been reported yet.

Our study aimed to estimate abundance and characterize *Psy* isolates from the water samples collected at the sites situated at the Danube River and its major tributaries in the territory of the Republic of Serbia and to determine their phenotypic and phylogenetic diversity. Isolates were identified by PCR reaction using species-specific primers. Their characterization was achieved through biochemical tests, hypersensitive reaction tests on *Pelargonium* plants, ice nucleation activity assay, motility assays, and *cts* gene-based phylogenetic analysis. The Danube and its tributaries were the aquatic habitats of choice, considering the Danube River Basin's potential and irrigation usage. This study provides a basis for a comprehensive pre-epidemic study of *P. syringae* in freshwaters in the context of irrigation, which should provide information on *Psy* community composition, abundance, and diversity.

Results

Abundance of *P. syringae* in the Danube River

Bacterial populations were evaluated for 15 sites along the Danube River Basin in Serbia (Fig. 1 and Additional file 1: Table S1). The total number of *Pseudomonads* spp. cultured on semi-selective media KBC ranged from 2.6×10^3 to 1.45×10^5 CFU/L (Additional file 1: Table S1). The total number of colonies cultured for each site was counted (D1—89, D2—85, D3—86, D4—145, D5—82, D6—95, D7—101, D8—102, D9—26, D10—91, D11—89, D12—69, T1—59, T2—52, T3—86 colonies). We treated all colonies as putative *Psy* isolates and performed PCR identification on all colonies grown on the filter. We performed a colony-PCR reaction using *Psy*-specific primers to detect *Psy* and reveal its abundance in the Danube. From the DNA of the isolates identified as *P. syringae*, 144-bp DNA fragments were amplified when

Psy-specific primers were used (Guilbaud et al. 2016). The specific band was amplified for 79 isolates belonging to the *P. syringae* species complex. Of 15 sampling sites, *Psy* was detected in ten, while none of the colonies tested were identified as *Psy* in the rest five sites (Fig. 1). The abundance of *P. syringae* varied from 1.0×10^2 to 1.0×10^4 CFU/L (Fig. 1 and Additional file 1: Table S1). The highest abundance was present at locality Tisa Titel (T1) with 1.2×10^4 CFU/L, and the lowest abundance was 1.0×10^2 CFU/L at localities Donji Milanovac (D8) and Tekija (D9).

Phylogenetic analysis

Further confirmation of the isolate identities was obtained by amplification and sequencing of the partial house-keeping citrate synthase (*cts*) gene. In 51 isolates, the *cts* gene had a significant percentage of identity




Fig. 1 Sampling locations on the Danube River (D1–D12) and its tributaries (T1–T3). Pie charts associated with sampling sites show the percentage share of detected *Pseudomonas syringae* phylogroups in the total number of isolates from that site. The *Psy* population sizes are indicated next to the pie charts and estimated in CFU per L. The black arrowhead shows the direction of the river flow

(98–100%) with *Psy* sequences from the NCBI nucleotide database, which were compared with query sequences using the blastn algorithm. These 51 isolates were subjected to further phenotypic and phylogenetic characterization. The remaining 28 isolates were excluded from further analysis due to low-quality reads of the partial *cts* gene and percentage identity below 98% similarity with *P. syringae*. Phylogenetic analysis based on partial

sequences of the *cts* gene (360 bp) involved 84 sequences (29 reference strains, four strains used as out-groups, and 51 sequences of the isolates from the Danube). The resulting phylogenetic analysis showed that isolates from the Danube belong to phylogroups (PG) 02 (70.6%), 07 (9.8%), 09 (9.8%), and 13 (9.8%) (Fig. 2). Groupings in PGs 02, 07, and 09 had bootstrap values of 77, 79, and 80, respectively, while the bootstrap value for PG13 was




Fig. 2 Phylogenetic tree of *Pseudomonas syringae* isolates based on the partial sequence of the *cts* gene. A phylogenetic relationship between 51 *Pseudomonas syringae* isolates from the Danube (BFD labeled isolates), reference strains representing 13 phylogroups, and strains used as out-groups (in bold) is shown. The tree was generated by the neighbor-joining method. Bootstrap values (expressed as a percentage of 1000 replications) are shown at branch nodes

93. The bootstrap value for clade-level classification in PG07 was 79 for clade 7a and 96 for clade 7b, which is significantly higher than the phylogroup level. Thirty-six out of 51 isolates were representatives of PG02, and these isolates were detected in each locality except Tekija (D9). Isolates BFD0003 and BFD0048 were placed into clades 2a and 2c, respectively, with high bootstrap values of 99 and 81. Five isolates were placed into PG07, PG09, and PG13. Isolate BFD0039 can be considered as a member of clade 9a with a maximum bootstrap value of 100. All tested isolates were distributed among these four *Psy* phylogroups, and there were no representatives of the remaining nine *Psy* phylogroups. Within all collected isolates from the Danube, we encountered 34 different *cts* gene sequences (haplotypes), while the dominant haplotype consisted of sequences belonging to PG02d. We compared the partial sequence of the *cts* gene (360 bp) of the strains detected in this study with those isolated from different diseased crops in Serbia and found 100% similarities between isolates from the Danube and isolates from diseased cherry plants (Ilicic et al. 2021) in Vojvodina Province. This haplotype (strains BFD0018, BFD0035, BFD0037, BFD0041, BFD0042, BFD0043, and BFD0049) was detected at five sampling sites, D1, D2, D3, T1, and T2 in the Danube River Basin (Fig. 1). Moreover, the same haplotype shared 100% identity with strains TAW79 and CC0170 isolated from water samples and cantaloupe in France (Berge et al. 2014; Guilbaud

et al. 2016). Another haplotype (isolates BFD0021 and BFD0036) showed 100% similarity to *Psy* strains isolated from diseased cherry plants which are also from Vojvodina Province in Serbia (Balaž et al. 2014), and to strain CC1435 isolated from an epilithic biofilm in France (Berge et al. 2014). In addition, we found high percent similarity (98.89%, four nucleotide differences) of strains BFD0017, BFD0019, BFD0023, and BFD0027 with *Psy* isolates from epidemics on pea and sugar beet in Serbia (Popović et al. 2015a; Nikolić et al. 2018).

Phenotypic characterization

Morphology of the colonies of each isolate grown on King's medium B was described after 24 h of incubation at 30°C (or after 48 h for slow-growing isolates). Colonies were predominantly small, white or creamy-white, circular, and flat, but within 51 isolates, we encountered orange-yellow colonies (three isolates) and brown pigmented (one isolate). Colony morphology description and the results of other tested phenotypic features are shown in Additional file 1: Table S1.

The pectinolytic activity was confirmed for three isolates (5.88%). Isolates positive for pectinolytic activity were representatives of PG07. The hypersensitive response was induced in *Pelargonium* plants by 35 *Psy* isolates (68.63%) (Fig. 3). 29 isolates of phylogroup 2 (56.86%), three isolates representing PG07 (60%), and




Fig. 3 Hypersensitive reaction assays. Top row, positive reactions induced by *Pseudomonas syringae* isolates from the Danube; bottom row, dH₂O used as a negative control and absence of hypersensitive response

three of PG09 (60%) could induce a hypersensitive response (Additional file 1: Table S1).

Ice nucleation activity

Three different types of ice nucleation activity depending on the temperature of ice formation were observed: warmer than -4°C (type I), -4°C to -7°C (type II), and colder than -7°C (type III) (Joly et al. 2013). Isolates could be classified into ice nucleators type I (7), type II (14), and type III (18), based solely on the average freezing temperatures (Fig. 4). Ice nucleation activity assays resulted in 39 active strains (all three tubes frozen), making up 76.47% of our collection (Fig. 5). INA active isolates from the Danube were from PG02 (94.44% of isolates), PG07 (80% of isolates), and PG09 (40% of isolates). The highest temperature that led to ice formation was -3°C for isolate BFD0052, based on an average value of triplicate tested. There were five more isolates with the capacity to form ice at -3°C (BFD0003, BFD0006, BFD0031, BFD0036, BFD0037), but not in each tested sample. All mentioned isolates that froze water at -3°C belong to PG02. Twelve isolates (BFD0003, BFD0006, BFD0008, BFD0018, BFD0024, BFD0026, BFD0029, BFD0031, BFD0043, BFD0045, BFD0050, BFD0056) cannot be precisely distinguished into categories as the temperature of freezing varies, and it is out of the assigned frame of categories.

Swimming and swarming motility

The swimming area was observed as a circular zone around the inoculation spot and measured using ImageJ

software (Additional file 2: Figure S1). Isolates BFD0030, BFD0040, and BFD0048 have shown poor swimming motility. Their growth around the inoculation spot reached 0.36, 0.46, and 0.1 cm^2 , respectively. The swimming area formed by the remaining 48 isolates was from 0.56 to 9.35 cm^2 . Isolate BFD0055 (representative of PG07) created the most extensive swimming area. *Pseudomonas aeruginosa* PAO1 was used as a positive control, and its swimming zone was 9.23 cm^2 . Regarding swarming motility, 25 isolates were swarming-positive. Examples of diverse movement patterns are shown in Additional file 2: Figure S2. The remaining 26 isolates have matched with negative control. All isolates from phylogroups 7 and 9 showed swarming motility. Isolates that were positive on swarming were also representatives of phylogroup 2 (41.67% of isolates) (Fig. 5). The smallest swarming area measured was 3.06 cm^2 , formed by isolate BFD0053. Isolates BFD0005 and BFD0001 from PG09 stood out with the biggest swarming areas 59.96 and 59.46 cm^2 , respectively.

Discussion

The presence of *P. syringae* in various environmental sources has been known for a long time, while the data about plant isolates still greatly surpasses the data about environmental isolates. Its ability to infect a wide range of hosts contributed to it becoming the best-studied model for understanding plant-microorganism interactions and pathogenicity. There is a great interest in *P. syringae* ecology, epidemiology, and evolution (Xin et al. 2018). They can persist in many habitats like streams, lakes, rivers,




Fig. 4 Average freezing temperature among ice-nucleation positive strains of *Pseudomonas syringae* from the Danube. INA positive strains are grouped into three types based on the average temperature of freezing (Type I $< -4^{\circ}\text{C}$, Type II -4°C to -7°C , and Type III $> -7^{\circ}\text{C}$)




Fig. 5 Phenotypic characterization of *Pseudomonas syringae* isolates from the Danube. **a** The percentage ratio of positive and negative results of phenotypic tests (individual results are provided in Additional file 1: Table S1). **b** Diagram showing the number of isolates sharing the common virulent traits. **c** Heatmap showing gradient in temperature of INA, motility area and capacity to degrade pectin and induce HR for isolates within each phylogroup. HR hypersensitive response, INA ice nucleation activity, PA pectinolytic activity

clouds, rain, and snow, and often lead an epiphytic life on disease-free plants (Morris et al. 2008). All these niches are potential reservoirs of a great diversity of strains and likely serve as sources of newly emerging pathogens (Dillon et al. 2019).

One of our goals was to determine *Psy* abundance in the Danube River. Our results show that the CFU/L of *Psy* ranges from 1.0×10^2 to 1.2×10^4 and that it was present in ten out of 15 sampled sites. Unfortunately, the Danube River in Serbia is affected by pollution from industrial development, agriculture, and poorly treated wastewater (Mănoiu and Crăciun 2021). Because organic

material accumulates strongly during filtration of concentrated water samples, we made serial dilutions of the samples, which in combination with the limited filter space possibly affected the range of *Psy* populations, but still allowed us to measure a *Psy* abundance of 10^4 CFU/L. Estimations of the population size of *Psy* isolated from different substrates like wild plants, snow, rain, epilithic biofilms, and lake and stream water can be found (Morris et al. 2008), but there is still insufficient information regarding *Psy* abundance in freshwater habitats. In lake and stream water, the abundance of *Psy* varied from 1.3×10^2 to 1.5×10^4 CFU/L (Morris et al.

2008). Also, in the headwaters of North America, Europe, and New Zealand rivers, the detected abundance of *Psy* was from 50 to 1.0×10^4 CFU/L (Morris et al. 2010). Population densities of *Psy* in the Durance River catchment, where it was monitored during four seasons over 2 years, were up to 1.0×10^5 CFU/L and its presence was confirmed in all sampling sites (Morris et al. 2022). Values of 1.0×10^5 CFU/L were reached during the winter seasons. Still, regarding spring seasons, population densities in the Durance River were lower, which coincides with the results obtained for the Danube River Basin.

The diversity of *Pseudomonas* species along the Danube River was recently assessed, resulting in 611 isolates identified by MALDI-TOF MS. Only two isolates were identified as a *P. syringae*, based on an additional *rpoD* gene sequence analysis (Mulet et al. 2020). The strains were isolated using an incubation temperature of 37°C. The overall diversity was probably much broader than acquired because the incubation and water temperatures during sampling (sampling was performed during late summer) were not optimal and not usually used to isolate environmental *Psy*. In fact, the impact of water temperature on the population size of *Psy* species in freshwater ecosystems has been shown recently, where *Psy* densities decreased with increasing water temperature, and it was the only factor among other aquatic parameters explaining population densities (Morris et al. 2022). Accordingly, the similarity of the occurrence of *Psy* populations in the lower part of the Durance River catchment, which coincides with the sampled sites on the Danube River, and in the Danube River in spring may be a consequence of similar temperature ranges in the same season. The aquatic parameters of the sampled Danube water showed a narrow range of measured values among different sites, and we did not find any correlation between *Psy* abundance and physicochemical parameters. Future seasonal monitoring would provide us with more information on the relationship between *Psy* abundance and variations in aquatic parameters in the Danube.

In the last decade, *Psy*-related disease epidemics on vegetables and fruits caused by *Psy* isolates from PG02 have been detected in the Serbian Danube River Basin. In particular, *Psy* has been identified as a causal agent of diseases in various crops in different parts of Vojvodina Province along the Danube-Tisa-Danube canal network. In particular, affected crops include oil pumpkins with disease incidence (DI) of 5–20% (Balaž et al. 2014), peas where DI reached 10–30% (Popović et al. 2015a), Swiss chard with DI 2–20% (Ignjatov et al. 2015), sugar beet with DI 0.1–40% (Stojšin et al. 2015), carrot, parsley and parsnip with DI 5–20% (Popović et al. 2015b). Among woody plants, its host was cherry, with DI up to 25% (Balaž et al. 2016; Ilicic et al. 2021; Iličić et al. 2022). The

highest DI of 80% was recorded in the blueberry orchard in Šabac, where 10% of the plants died (Zlatković et al. 2022). All these reports represent disease occurrence at sites surrounded by the Danube River and the Danube-Tisa-Danube Canal. We reasoned that these freshwater bodies could be a possible source of *Psy* inoculum and consequently a risk for crops in this area. The concordance of *cts* sequences from two haplotypes of *Psy* strains isolated from the Danube River and the causal agents of the reported disease on cherry plants suggests the presence of mixed environmental and agricultural populations and the putative risk of the Danube irrigation system for future disease occurrence. However, long-term monitoring of the Danube River Basin is needed to assess the extent of this risk.

The considerable genotypic and phenotypic heterogeneity of *Psy* species makes it difficult to access the full diversity of this group during isolation (Berge et al. 2014). Phenotypic comparisons with some previously described reference strains may result in the loss of a substantial number of *Psy* group members. A PCR-based method for detecting *Pseudomonas syringae* using *Psy*-specific primers (Guilbaud et al. 2016) is much more comprehensive than isolation based on phenotypic characteristics. It allowed us to cover the entire *Psy* group and isolate different strains that show great phenotypic diversity within the *Psy* collection from the Danube River. In addition, the *cts* gene has previously been found to be one of the most reliable genes that can be used for determining the phylogenetic affiliation of *Psy* strains (Berge et al. 2014). We observed the greatest abundance of *Psy* isolates belonging to PG02 and the highest phylogenetic and phenotypic diversity within this phylogroup among the whole *Psy* isolate collection from the Danube. PG02 is the most ubiquitous group for which numerous isolates have been found in environmental habitats, including water sources (Berge et al. 2014). Additionally, strains from rivers that belong to PG02 are more aggressive on cantaloupe seedlings than strains in other phylogroups (Berge et al. 2014). Isolates from PG02 are known as potential ice nucleators (Pietsch et al. 2017), and previous estimations show that 85% of isolated strains from PG02 were INA-positive (Morris et al. 2010). Members of PG02 isolated from freshwaters have been linked on the basis of their *cts* gene sequences to strains characterized as worldwide causative agents of reported epidemics in apricots, melons, squash, and sugar beets (Morris et al. 2022). Morris et al. (2022) identified the haplotype DD.1 (PG02b) as the most abundant in their *Psy* strain collection from the Durance River catchment, whose partial sequence of the *cts* gene has 100% similarity to sequences of *Psy* previously isolated from numerous disease epidemics worldwide, including sugar beet

epidemics in Serbia. Although we detected 34 different haplotypes based on the *cts* gene in the strain collection from the Danube River Basin, we did not find 100% similarity to the DD.1 haplotype (the highest similarity was 98.89%, four nucleotide differences, in strains BFD0017, BFD0019, BFD0023, and BFD0027). The representative of PG02c, isolate BFD0048, did not elicit a hypersensitive response in the *Pelargonium* plant. Strains from clade 2c are known to be members of the *Pseudomonas congelans* species (Behrendt et al. 2003). Although they have been primarily described as non-pathogenic (Dillon et al. 2019), their involvement in the development of diseases in citrus plants in Tunisia has been reported (Oueslati et al. 2020). In addition, the pathogenicity of *P. congelans* isolated from the phyllosphere of Serbian autochthonous plum cultivars was reported with a severity index of 44% on the leaves (Janakiev et al. 2020).

P. syringae isolates of other phylogroups detected in the Danube River (PG07, 09, and 13) were not often detected as pathogens of diseases in crops in Serbia. To date, there was one initial report of *P. viridiflava* (PG07) in the southern part of Serbia (outside the Danube River Basin) with a disease incidence of 10–25% on tomato plants (Popović et al. 2015c). Strains from this phylogroup have also been previously isolated from water habitats, and they are known as *P. viridiflava* representatives of the *Psy* species complex (Lipps and Samac 2022). Their prominent feature is the production of the enzyme pectate lyase, which is used to degrade pectin in plant cell walls as one of the main virulence factors (Lipps and Samac 2022). Moreover, by an investigation from the beginning of the twenty-first century until 2015, *P. viridiflava* was responsible for 18% of all disease outbreaks caused by the *Psy* species complex (Lamichhane et al. 2015). PG09 strains have been reported exclusively from aquatic habitats and are considered to be well adapted to the environment, while PG13 is also widespread but mostly found on non-plant substrates (Berge et al. 2014).

To get additional insight into the colonization capabilities of *Psy* strains from the Danube which could be considered as an important pathogenic feature, we examined their motility. The ability to move toward nutrients, attach, and penetrate host tissues is an excellent advantage for a pathogenic lifestyle (Colin et al. 2021). Examined forms of motility, swimming and swarming, are flagellum-mediated and depend on environmental conditions (Markel et al. 2018). While swarming is locomotion on a semi-solid surface, swimming occurs in the liquid phase (Jose and Singh 2020). A connection between swarming motility and host range extent was detected earlier, and strains with a tendency to swarm in the first 24 h have been shown to have a broader host range than non-swarming ones (Morris et al. 2019). In

our collection, we encountered numerous strains that stand out with their ability to swim and swarm in the given conditions, including strains from PG02, which, according to the data available so far, pose the greatest threat to plants in the Danube River Basin. That suggests that among the *Psy* isolates from the Danube, those with the largest motility areas can pose a threat to plant health because this trait makes them a potential pathogen with increased virulence.

By comparing sequenced genomes of environmental and crop-pathogenic strains and identifying genes with key roles in disease emergence, the hypothesis that crop-pathogenic *P. syringae* diverged from a pre-existing population that was present in the environment before the development of modern agriculture was suggested (Monteil et al. 2016). A significant genetic similarity between strains from nonagricultural habitats and those isolated from disease epidemics was shown (Morris et al. 2008; Monteil et al. 2013). Mixing populations from agricultural and environmental sources is highly possible, bearing in mind that they are genetically almost indistinguishable (Bartoli et al. 2015; Monteil et al. 2016). These data increase the importance of studies that provide insight into the diversity of populations outside of agroecosystems. Regarding quite frequent reports of plant diseases caused by strains from the *Psy* complex (especially PG02), high diversity among *Psy* strains from the Danube River, and extensive use of Danube freshwater for crop irrigation purposes, future studies should be focused on monitoring seasonal population changes and determination of pathogenicity and the host range of the isolates in order to assess the potential risk for irrigation of certain crops and even predict potential epidemics.

Conclusions

This study represents the first report on the abundance and diversity of the plant pathogen *P. syringae* in the Danube River Basin. We revealed the abundance of *Psy* in the Danube River, and the presence of four phylogroups of *Psy*, including isolates with confirmed putative virulent traits such as ice nucleation activity, flagellum-dependent motility, and pectinolytic activity. It remains to be investigated whether the Danube River Basin is a possible source of pathogen inoculum for irrigated crops.

Methods

Sample sites and collection

Samples were collected from 11 to 14th May 2021 from 15 sites (Fig. 1). Sites D1–D12 were situated on the Danube River, while sites T1–T3 were located at the Danube tributaries. In the investigated stretch, the Danube has characteristics of a large lowland river. Except for Site T3 (the River Pek), all sites are heavily modified water bodies

affected by the Iron Gate dams. The construction of the Iron Gate dams resulted in hydromorphological changes in this part of the river and the formation of reservoirs with reduced sediment flux, increased sediment deposition, and slowed river flow (Vuković et al. 2014). All samples were collected approximately 3 m from the shore at a 30 cm depth below the surface in clean, sterile 500 mL bottles. Samples were immediately stored at 4°C and transported to the laboratory in cooling boxes (within 3 h). In-field measurements (temperature, pH, conductivity, and dissolved oxygen levels) were performed using a multi-parameter probe (WTW/Xylem Analytics, Germany). Data on measurements are provided in the Additional file 1: Table S1.

Sample processing and PCR detection of *P. syringae*

Water samples were serially diluted in sterile distilled water (tenfold, 100-fold, and 1000-fold) in the final volume of 100 mL and filtered. Membrane filtration was done using a mixed cellulose esters filter with a pore diameter of 0.45 µm and a filter diameter of 47 mm (Millipore, France). Filters were transferred face up on the surface of KBC media (Mohan and Schaad 1987), selective for the growth of *Pseudomonas* species, and incubated for 3–5 days at room temperature. Since we filtered three different dilutions of sampled river water for abundance estimation and *Psy* identification, we further processed dilutions with a countable number of colonies (100 colonies or less per filter for tenfold dilutions (47 mm diameter filter), except for the D4 site with 145 colonies in 100-fold dilution). In total, 1257 grown colonies from the filters were subjected to a *Psy*-specific PCR reaction to detect *Pseudomonas syringae* species using primers named *Psy_F* and *Psy_R*, designed by Guilbaud et al. (2016). PCR reactions were conducted in MiniAmp™ Thermal Cycler (Thermo Fischer Scientific, Waltham, Massachusetts, USA). PCR mixture and conditions used were also described by Guilbaud et al. (2016). In brief, PCR reactions were conducted in a final volume of 25 µL with each mix containing 14 µL of PCR water (Thermo Fischer Scientific, Waltham, Massachusetts, USA), 5 µL of 10× KAPA Taq buffer (KAPA Biosystems Inc, USA), 1.5 µL of MgCl₂ (25 mM), 0.3 µL of dNTP mix (2.5 mM per nucleotide), 0.2 µL of KAPA Taq Polymerase (5 U/µL, KAPA Biosystems Inc, USA) and 1 µL of each *Psy*-specific primer (10 mM). Bacterial cell material from single colonies was picked up with a sterile 10 µL pipette tip and transferred directly into PCR tubes with a PCR mixture. PCR reactions were conducted with an initial denaturation step for 5 min at 96°C, followed by 30 cycles at 94°C for 30 s, 61°C for 30 s, 72°C for 30 s, and final elongation for 10 min at 72°C. PCR products were visualized on 1% agarose gel containing an aqueous solution of

10 mg/mL ethidium-bromide (SERVA, Germany). Electrophoresis was run in Tris–borate-EDTA buffer (5.4 g Tris; 2.75 g Boric acid; 4 mL 0.5 M pH 8 EDTA; dH₂O to 1 L) at a constant voltage of 90 V and 300 mA amperage for 60 min. *Psy*-PCR positive strains were stocked in LB medium with 20% glycerol and stored at –80°C.

DNA extraction

For total genomic DNA extraction, the modified CTAB protocol proposed by Le Marrec et al. (2000) was used. Single bacterial colonies of each isolate were resuspended in a mix of 567 µL of TE buffer (pH 7.6; 10 mM Tris; 1 mM EDTA), 30 µL of 10% SDS, and 3 µL 20 mg/mL of proteinase K. After incubation at 37°C for 30 min, 100 µL of 5 M NaCl and 300 µL of 3% CTAB+PVP buffer (pH 8; 3% CTAB; 1 M Tris; 1.4 M NaCl; 20 mM EDTA; 3% PVP) was added, vortexed, and incubated on the heating block at 65°C for 20 min. In the next step, the DNA was purified with 800 µL of chloroform, vortexed thoroughly, and centrifugated for 10 min at 8000 g. Supernatant in the upper phase was transferred to a new tube and mixed with 3 M sodium acetate (pH 5) in a volume ratio of 1:10. An equal volume of ice-cold isopropanol (~750 µL) was added and the mix was centrifugated for 15 min at 8000 g. The liquid phase was discarded, and the precipitate was washed with 1 mL of ice-cold 96% ethanol and centrifugated for 10 min at 8000 g. The liquid phase was discarded again, and the precipitate was dried for 30 min at 37°C and resuspended in 50 µL of TE buffer with RNase mix added in a final concentration of 0.2 mg/mL. The isolated genomic DNA was incubated for 15 min at 37°C and stored at –20°C.

Phylogenetic analysis

To further confirm *P. syringae* identity and their phylogeny, a partial sequence of citrate synthase house-keeping gene (*cts*) was amplified and sequenced. DNA amplification and sequencing were done with primers described previously (Morris et al. 2010). The PCR reaction was done in a final volume of 50 µL containing 25 µL of DreamTaq™ Green PCR Master Mix 2× and 21 µL of PCR water (Thermo Fischer Scientific, Waltham, Massachusetts, USA), 1 µL of each primer (10 µmol) and 2 µL of template DNA. PCR reactions were conducted with initial denaturation for 5 min at 96°C, followed by 35 cycles at 94°C for 30 s, 62°C for 1 min 30 s, 72°C for 2 min, and final elongation for 7 min at 72°C. PCR products were purified with GeneJET PCR Purification Kit (Thermo Fischer Scientific, Waltham, Massachusetts, USA) and Zymoclean™ Gel DNA Recovery Kit (Zymo Research, USA). PCR products were visualized on 1% agarose gel containing an aqueous solution of 10 mg/mL ethidium-bromide (SERVA, Germany). Electrophoresis

was run in Tris–borate–EDTA buffer (5.4 g Tris; 2.75 g Boric acid; 4 mL 0.5 M pH 8 EDTA; dH₂O to 1 L) at a constant voltage of 90 V and 300 mA amperage for 90 min for partial *cts* gene (citrate synthase gene) products. Agarose gels were visualized with a transilluminator (LKB, Transilluminator 2011 Microvue UV Light, Sweden). All amplicons were sequenced by Eurofins Genomics GmbH (Wien, Austria) using the sequencing primer described previously (Morris et al. 2010). Sequences were processed using the FinchTV v. 1.4.0 software package (Geospiza Inc.). Sequencing was repeated up to three times for certain samples to avoid sequencing errors and obtain more reliable results. Sequences that had low quality or less than 98% similarity to *Psy* were excluded from phylogenetic analysis. Sequences of representative *P. syringae* strains were retrieved from Berge et al. (2014) and included in the phylogenetic analysis to determine the phylogenetic grouping of *P. syringae* strains isolated from the Danube. Selected reference strains included one representative of each phylogroup and clades within phylogroups. Among reference strains, there was chosen one strain representing PG03 (LYR0002), PG04 (CC1513), PG05 (PmaES4326), PG06 (CFBP2067), PG08 (GAW0203), and PG11 (CFBP4407), two strains representing PG01 (DC3000—clade 1a; CC1416—1b), PG07 (CC1582—7a; FMU107—7b), PG12 (GAW0112—12a; GAW0113—12b), and PG13 (CCE0915—13a; CCV0567—13b), three from PG09 (CC1532, TA0006, CMW0020—clades from a to c, respectively), five from PG02 (PsyCit7, H5E1, PSy642, CC1470, USA0035—clades a to e, respectively) and seven strains representing PG10 (CC1583, TA0019, CC1586, CCE0100, USA0032, CCE0153, CCV01213—clades a to g, respectively). Reference strains *P. graminis*, *P. rhizosphaerae*, *P. protegens*, and *P. aeruginosa* were included as out-groups, as used by Berge et al. (2014). The tree was rooted on the *P. aeruginosa* PAO1 strain. The *cts* gene sequences were aligned using the Muscle program integrated into the Mega 11 software and used to create a dendrogram in Mega 11, inferring evolutionary history using the neighbor-joining method. The percentage of replicate trees where associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches. Sequences from this research were deposited in the NCBI database under the following accession numbers: OP620590–OP620592, OP620594–OP620602, OP620606–OP620613, OP620615–OP620616, OP620618–OP620626, OP620628–OP620635, OP620637–OP620646, and OP620648–OP620649. The *cts* gene partial sequences used for comparison among all collected isolates from the Danube were subsequently blasted using the NCBI BLASTn tool in order to assess similarities between sequences and distinguish haplotypes.

Phenotypic characterization

Phenotypic characterization of 51 *Psy* isolates previously confirmed by sequencing of partial *cts* gene included a description of colony morphology, pectinolytic activity, induction of a hypersensitive reaction, ice nucleation activity assay, swimming, and swarming assays.

Besides the isolates from the Danube, other bacterial strains used in phenotypic characterization as controls were: *P. aeruginosa* PAO1, *P. syringae* pv. *aptata* P5, and *Pectobacterium carotovorum* 2811. Strains used in this study were cultivated as overnight cultures in KB broth (30°C with shaking at 180 rpm, except *P. aeruginosa* PAO1 that was incubated at 37°C) or streaked on a solid KB (24 h incubation at 30°C). Depending on the phenotypic test, adjusting optical density was done in fresh KB broth to OD₆₀₀ ~ 0.2 (approximately 1.0 × 10⁸ CFU/mL) and OD₆₀₀ ~ 0.3 to 0.4 (approximately 2.0 × 10⁸ CFU/mL). Optical density adjustment was made based on the growth curves of two randomly selected isolates from the Danube River. In brief, overnight cultures of the isolates BFD0012 and BFD0052 were diluted 50-fold in 50 mL of fresh KB medium and incubated at 30°C with shaking at 180 rpm. OD₆₀₀ was measured every 45 min. In parallel, samples from the incubated culture were taken at a given checkpoint, serially diluted in 0.01 M MgSO₄, and 0.1 mL of dilutions was poured and spread plated onto solid KB medium. After incubation at 30°C for 24 h, the CFU/mL was calculated. Obtained values of optical density and CFU/mL for each checkpoint were utilized for constructing a standardization curve for easy use of optical density to approximate bacterial cell density (Additional file 2: Figure S3). For both isolates, there are visible lag phases and gradual entry in the exponential phase of growth which is followed by the beginning of the stationary phase. For BFD0012, cell density reaches 1.0 × 10⁸ CFU/mL at OD₆₀₀ = 0.212 and 2.0 × 10⁸ at OD₆₀₀ = 0.318. For BFD0052, cell density was 1.0 × 10⁸ CFU/mL at OD₆₀₀ = 0.156 and 2.0 × 10⁸ at OD₆₀₀ = 0.284.

Pectinolytic activity

The pectinolytic activity was tested using 1 cm thick potato slices placed in a sterile Petri dish with filter paper soaked in sterile water. 100 µL of bacterial suspensions with OD₆₀₀ adjusted to ~ 0.2 were inoculated into indentations made in potato slices with a sterile 1 mL pipette tip. Each isolate was tested in triplicate. Results were observed after 48 h of incubation at room temperature. Softened, brown tissue around the indentations was a

sign of a positive reaction. The positive control used for this test was the *P. carotovorum* 2811 strain.

Hypersensitive reaction

The hypersensitive reaction was tested on *Pelargonium* plants. Bacterial suspensions were adjusted to $OD_{600} \sim 0.2$, centrifugated for 10 min at 2000 g, and resuspended in sterile distilled water. Using a needleless medical syringe, *Pelargonium* leaves were inoculated with bacterial suspensions on the abaxial side between two lateral veins. Each isolate was tested in triplicate. Sterile distilled water was used as a negative control. Results were assessed 24–48 h and 7 days post-inoculation.

Ice nucleation activity assay

Ice nucleation activity was tested using a CH-100 Cooling Dry Block (Biosan, Riga, Latvia) with a protocol modified and optimized according to Nemecek-Marshall et al. (1993) and Joly et al. (2013). From overnight cultures, 2 mL volume was exposed to a temperature of 15°C for 1 h in the stationary growth phase to induce expression of ice nucleation phenotype. After induction, the optical density of isolates was adjusted to $OD_{600} \sim 0.2$ in the final volume of 1 mL in a fresh KB medium. The bacterial suspensions were centrifugated for 10 min at 2000 g, and the precipitate was resuspended in the same volume of sterile dH_2O . 20 μ L of prepared bacterial suspensions, which contain 2.0×10^6 CFU/mL, was distributed to 0.2 mL PCR tubes, each isolate in triplicate (three tubes). PCR tubes were placed in a Cooling Dry Block, and isolates were exposed to decreasing temperatures from -2°C to -10°C in a decline of 1°C . Samples were checked for ice formation every 5 min (freeze check was performed with sterile tips), an interval of exposition to a given temperature. Sterile dH_2O was used as a negative control. If ice was present in two of three tubes in a given temperature range, the isolate was considered INA+. The temperature at which each of the tubes froze was recorded.

Swimming and swarming assays

For motility assays, the optical density of bacterial suspensions was adjusted to $OD_{600} \sim 0.3$. Swimming media was prepared as 50% KB containing 0.25% agar and swarming media as undiluted KB containing 0.4% agar, as described in Hockett et al. (2013). A swimming assay was performed by stabbing bacterial cell suspension in the center of a swimming plate with a sterile 10 μ L pipette tip. For the swarming assay, 3 μ L aliquots were inoculated onto the center of a swarming plate.

The swimming and swarming motilities were observed 20–24 h after incubation at room temperature. ImageJ software was used to measure swimming and swarming areas (Schneider et al. 2012). *P. aeruginosa* PAO1 strain was used as a positive control for swimming motility (Yang et al. 2018). The negative control for the swarming motility assay was *P. syringae* pv. *aptata* P5 strain (Morris et al. 2019).

Abbreviations

cts	Citrate-synthase
INA	Ice nucleation activity
PG	Phylogroup
Psy	<i>Pseudomonas syringae</i>

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s42483-023-00174-0>.

Additional file 1. Table S1. Abundance and phenotypic characterization of *Pseudomonas syringae* collection from the Danube River and physico-chemical parameters of freshwater samples measured on-site. P, pectinolytic activity; HR, hypersensitive response on *Pelargonium* plants; INA, ice nucleation activity; Swimming and swarming area measured in ImageJ and expressed in cm^2

Additional file 2. Figure S1. Swimming area (cm^2) of *Pseudomonas syringae* isolates from the Danube measured with ImageJ software. BFD0026, BFD0040, and BFD0059: selected isolates showing gradation in swimming motility; PAO1: *Pseudomonas aeruginosa* was used as a positive control. **Figure S2.** Swarming area (cm^2) of *Pseudomonas syringae* isolates from the Danube measured with ImageJ software. BFD0008, BFD0024, BFD0029, BFD0039, BFD0042, BFD0052, BFD0057, BFD0060: selected isolates showing swarming motility; C: *Pseudomonas syringae* pv. *aptata* P5 was used as a negative control. **Figure S3.** Growth curves of two *Pseudomonas syringae* isolates from the Danube River. **a** *Pseudomonas syringae* BFD0012 OD_{600} and CFU/mL growth curve and standard growth curve. **b** *Pseudomonas syringae* BFD0052 OD_{600} and CFU/mL growth curve and standard growth curve.

Acknowledgements

We would like to acknowledge researchers from the Department of Hydroecology and Water Protection of the Institute for Biological Research 'Siniša Stanković' for collaboration and support in sample collection for this research.

Authors' contributions

IN, TB, and SS designed the research. SK provided water samples for the research and initial sample processing. MA and IR carried out the experiments. MA wrote the first draft of the manuscript, and IR, OM, SK, TB, SS, and IN contributed to the final version. All authors read and approved the submitted version of the manuscript.

Funding

This work was supported by the Ministry of Science, Technological Development and Innovations of the Republic of Serbia, grants No. 451-03-47/2023-01/200178 (Faculty of Biology) and 451-03-47/2023-01/200007 (Institute for Biological Research 'Siniša Stanković').

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 17 November 2022 Accepted: 10 April 2023

Published online: 19 May 2023

References

- Balaž J, Iličić R, Maširević S, Jošić D, Kojić S. First report of *Pseudomonas syringae* pv. *syringae* causing bacterial leaf spots of oil pumpkin (*Cucurbita pepo*) in Serbia. *Plant Dis.* 2014;98(5):684. <https://doi.org/10.1094/PDIS-07-13-0714-PDN>.
- Balaž J, Iličić R, Ognjanov V, Ivanović Ž, Popović T. Etiology of bacterial canker on young sweet cherry trees in Serbia. *J Plant Pathol.* 2016;98:285–94.
- Banciu AR, Ionica DL, Vaideanu MA, Radulescu DM, Nita-Lazar M, Covaliu CI. The occurrence of potentially pathogenic and antibiotic resistant Gram-negative bacteria isolated from the Danube Delta ecosystem. *Sustainability.* 2021;13(7):3955. <https://doi.org/10.3390/su13073955>.
- Baroudy F, Habib W, Tanos G, Gerges E, Saab C, Choueiri E, et al. Long-distance spread of *Verticillium dahliae* through rivers and irrigation systems. *Plant Dis.* 2018;102(8):1559–65. <https://doi.org/10.1094/PDIS-08-17-1189-RE>.
- Behrendt U, Ulrich A, Schumann P. Fluorescent pseudomonads associated with the phyllosphere of grasses; *Pseudomonas trivialis* sp. Nov., *Pseudomonas poae* sp. nov. and *Pseudomonas congelans* sp. Nov. *Int J Syst Evol Microbiol.* 2003;53(5):1461–9. <https://doi.org/10.1099/ijs.0.02567-0>.
- Berge O, Monteil CL, Bartoli C, Chandeysson C, Guilbaud C, Sands DC, et al. A user's guide to a data base of the diversity of *Pseudomonas syringae* and its application to classifying strains in this phylogenetic complex. *PLoS ONE.* 2014;9(9):e105547. <https://doi.org/10.1371/journal.pone.0105547>.
- Bartoli C, Lamichhane JR, Berge O, Guilbaud C, Varvaro L, Balestra GM, et al. A framework to gauge the epidemic potential of plant pathogens in environmental reservoirs: the example of kiwifruit canker. *Mol Plant Pathol.* 2015;16(2):137–49. <https://doi.org/10.1111/mpp.12167>.
- Colin R, Ni B, Laganenka L, Sourjik V. Multiple functions of flagellar motility and chemotaxis in bacterial physiology. *FEMS Microbiol Rev.* 2021;45(6):fuab038. <https://doi.org/10.1093/femsre/fuab038>.
- Dávid A, Madudová E. The Danube river and its importance on the Danube countries in cargo transport. *Transp Res Proc.* 2019;40:1010–6. <https://doi.org/10.1016/j.trpro.2019.07.141>.
- De Araujo GG, Rodrigues F, Gonçalves FL, Galante D. Survival and ice nucleation activity of *Pseudomonas syringae* strains exposed to simulated high-altitude atmospheric conditions. *Sci Rep.* 2019;9(1):7768. <https://doi.org/10.1038/s41598-019-44283-3>.
- Dillon MM, Thakur S, Almeida RN, Wang PW, Weir BS, Guttman DS. Recombination of ecologically and evolutionarily significant loci maintains genetic cohesion in the *Pseudomonas syringae* species complex. *Genome Biol.* 2019;20(1):1–28. <https://doi.org/10.1186/s13059-018-1606-y>.
- Donati I, Cellini A, Sangiorgio D, Vanneste JL, Scortichini M, Balestra GM, et al. *Pseudomonas syringae* pv. *actinidiae*: ecology, infection dynamics and disease epidemiology. *Microb Ecol.* 2020;80(1):81–102. <https://doi.org/10.1007/s00248-019-01459-8>.
- Faye P, Bertrand C, Pédrón J, Barny MA. Draft genomes of “*Pectobacterium peruviansense*” strains isolated from fresh water in France. *Stand Genom Sci.* 2018;13(1):1–8. <https://doi.org/10.1186/s40793-018-0332-0>.
- Fayette J, Jones JB, Pernezny K, Roberts PD, Raid R. Survival of *Xanthomonas campestris* pv. *vitiensis* on lettuce in crop debris, irrigation water, and weeds in south Florida. *Eur J Plant Pathol.* 2018;151(2):341–53. <https://doi.org/10.1007/s10658-017-1377-4>.
- Frick C, Vierheilig J, Nadiotis-Tsaka T, Ixenmaier S, Linke R, Reischer GH, et al. Elucidating fecal pollution patterns in alluvial water resources by linking standard fecal indicator bacteria to river connectivity and genetic microbial source tracking. *Water Res.* 2020;184:116132. <https://doi.org/10.1016/j.watres.2020.116132>.
- Guilbaud C, Morris CE, Barakat M, Ortet P, Berge O. Isolation and identification of *Pseudomonas syringae* facilitated by a PCR targeting the whole *P. syringae* group. *FEMS Microbiol Ecol.* 2016;92(1):fv146. <https://doi.org/10.1093/femsec/fv146>.
- Gutiérrez-Barranquero JA, Cazorla FM, De Vicente A. *Pseudomonas syringae* pv. *syringae* associated with mango trees, a particular pathogen within the “hodgepodge” of the *Pseudomonas syringae* complex. *Front Plant Sci.* 2019;10:570. <https://doi.org/10.3389/fpls.2019.00570>.
- Hockett KL, Burch AY, Lindow SE. Thermo-regulation of genes mediating motility and plant interactions in *Pseudomonas syringae*. *PLoS ONE.* 2013;8(3):e59580. <https://doi.org/10.1371/journal.pone.0059580>.
- Hugouvieux-Cotte-Pattat N, Jacot-des-Combes C, Briolay J. *Dickeya lacustris* sp. Nov., a water-living pectinolytic bacterium isolated from lakes in France. *Int J Syst Evol Microbiol.* 2019;69(3):721–6. <https://doi.org/10.1099/ijsem.0.003208>.
- Ignjatov M, Gvozdanović-Varga J, Milošević D, Nikolić Z, Ivanović Ž, Popović T. First report of bacterial leaf spot of chard (*Beta vulgaris* subsp. *cicla*) caused by *Pseudomonas syringae* pv. *syringae* in Serbia. *Plant Dis.* 2015;99(5):723. <https://doi.org/10.1094/PDIS-10-14-1097-PDN>.
- Iličić R, Popović T, Marković S, Jelušić A, Bagi F, Vljajić S, et al. Genetic diversity of *Pseudomonas syringae* pv. *syringae* isolated from sweet cherry in southern and northern regions in Serbia. *Genetika-Belgrade.* 2021;53(1):247–62. <https://doi.org/10.2298/GENSR2101247I>.
- Iličić R, Jelušić A, Marković S, Barać G, Bagi F, Popović T. *Pseudomonas cerasi*, the new wild cherry pathogen in Serbia and the potential use of recG helicase in bacterial identification. *Ann Appl Biol.* 2022;180(1):140–50. <https://doi.org/10.1111/aab.12717>.
- Janakiev T, Unković N, Dimkić I, Ljaljević-Grbić M, Stević T, Stanković S, et al. Susceptibility of Serbian plum cultivars to indigenous bacterial and *Monilinia laxa* isolates. *Bot Serb.* 2020;44(2):203–10. <https://doi.org/10.2298/BOTSERB2002203J>.
- Joly M, Attard E, Sancelme M, Deguillaume L, Guilbaud C, Morris CE, et al. Ice nucleation activity of bacteria isolated from cloud water. *Atmos Environ.* 2013;70:392–400. <https://doi.org/10.1016/j.atmosenv.2013.01.027>.
- Jose R, Singh V. Swarming in bacteria: a tale of plasticity in motility behavior. *J Indian Inst Sci.* 2020;100(3):515–24. <https://doi.org/10.1007/s41745-020-00177-2>.
- Karimi B, Nosrati R, Fazly Bazzaz BS, Mirpour M, Malboobi M, Owlia P. A comparative evaluation of freezing criteria and molecular characterization of epiphytic ice-nucleating (Ice+) and non-ice-nucleating (Ice-) *Pseudomonas syringae* and *Pseudomonas fluorescens*. *J Plant Pathol.* 2020;102(1):169–78. <https://doi.org/10.1007/s42161-019-00402-7>.
- Kirschner AK, Reischer GH, Jakwerth S, Savio D, Ixenmaier S, Toth E, et al. Multiparametric monitoring of microbial faecal pollution reveals the dominance of human contamination along the whole Danube River. *Water Res.* 2017;124:543–55. <https://doi.org/10.1016/j.watres.2017.07.052>.
- Kurilić SM, Staletović N, Abulsba KS, Dagali KS. Possibility of the application of Danube, Sava and Morava River area shallow aquifer groundwater (Serbia) in the irrigation purposes. *Water Resour.* 2019;46(6):861–70. <https://doi.org/10.1134/S0097807819060162>.
- Lamichhane JR, Messéan A, Morris CE. Insights into epidemiology and control of diseases of annual plants caused by the *Pseudomonas syringae* species complex. *J Gen Plant Pathol.* 2015;81(5):331–50. <https://doi.org/10.1007/s10327-015-0605-z>.
- Laurila J, Ahola V, Lehtinen A, Joutsjoki T, Hannukkala A, Rahkonen A, et al. Characterization of *Dickeya* strains isolated from potato and river water samples in Finland. *Eur J Plant Pathol.* 2008;122(2):213–25. <https://doi.org/10.1007/s10658-008-9274-5>.
- Le Marrec C, Hyronimus B, Bressollier P, Verneuil B, Urdaci MC. Biochemical and genetic characterization of coagulin, a new antilisterial bacteriocin in the pediocin family of bacteriocins, produced by *Bacillus coagulans* I4. *Appl Environ Microbiol.* 2000;66(12):5213–20. <https://doi.org/10.1128/AEM.66.12.5213-5220.2000>.
- Lipps SM, Samac DA. *Pseudomonas viridiflava*: an internal outsider of the *Pseudomonas syringae* species complex. *Mol Plant Pathol.* 2022;23(1):3–15. <https://doi.org/10.1111/mpp.13133>.

- Maki LR, Galyan EL, Chang-Chien MM, Caldwell DR. Ice nucleation induced by *Pseudomonas syringae*. *Appl Microbiol*. 1974;28(3):456–9. <https://doi.org/10.1128/am.28.3.456-459.1974>.
- Mănoiu VM, Crăciun AI. Danube river water quality trends: a qualitative review based on the open access web of science database. *Ecohydrol Hydrobiol*. 2021;24(4):613–28. <https://doi.org/10.1016/j.ecohyd.2021.08.002>.
- Markel E, Dalenberg H, Monteil CL, Vinatzer BA, Swingle B. An AlgU-regulated antisense transcript encoded within the *Pseudomonas syringae* fleQ gene has a positive effect on motility. *J Bacteriol*. 2018;200(7):e00576–e617. <https://doi.org/10.1128/JB.00576-17>.
- Mohan SK, Schaad NW. An improved agar plating assay for detecting *Pseudomonas syringae* pv. *syringae* and *P. s.* pv. *phaseolicola* in contaminated bean seed. *Phytopathology*. 1987;77(10):1390–5.
- Monteil CL, Cai R, Liu H, Mehan L, Lontop ME, Leman S, Studholme DJ, et al. Nonagricultural reservoirs contribute to emergence and evolution of *Pseudomonas syringae* crop pathogens. *New Phytol*. 2013;199(3):800–11. <https://doi.org/10.1111/nph.12316>.
- Monteil CL, Yahara K, Studholme DJ, Mageiros L, Méric G, Swingle B, et al. Population-genomic insights into emergence, crop adaptation and dissemination of *Pseudomonas syringae* pathogens. *Microb Genom*. 2016. <https://doi.org/10.1099/mgen.0.000089>.
- Morris CE, Sands DC, Vinatzer BA, Glaux C, Guilbaud C, Buffiere A, et al. The life history of the plant pathogen *Pseudomonas syringae* is linked to the water cycle. *ISME J*. 2008;2(3):321–34. <https://doi.org/10.1038/ismej.2007.113>.
- Morris CE, Bardin M, Kinkel LL, Moury B, Nicot PC, Sands DC. Expanding the paradigms of plant pathogen life history and evolution of parasitic fitness beyond agricultural boundaries. *PLoS Pathog*. 2009;5(12):e1000693. <https://doi.org/10.1371/journal.ppat.1000693>.
- Morris CE, Sands DC, Vanneste JL, Montary J, Oakley B, Guilbaud C, et al. Inferring the evolutionary history of the plant pathogen *Pseudomonas syringae* from its biogeography in headwaters of rivers in North America, Europe, and New Zealand. *Mbio*. 2010;1(3):e00107–10. <https://doi.org/10.1128/mBio.00107-10>.
- Morris CE, Lamichhane JR, Nikolić I, Stanković S, Moury B. The overlapping continuum of host range among strains in the *Pseudomonas syringae* complex. *Phytopathol Res*. 2019;1:1–6. <https://doi.org/10.1186/s42483-018-0010-6>.
- Morris CE, Lacroix C, Chandeysson C, Guilbaud C, Monteil C, Piry S, et al. Comparative seasonal abundance and diversity of populations of the *Pseudomonas syringae* and Soft Rot *Pectobacteriaceae* species complexes throughout the Durance River catchment from its French Alps sources to its delta. *bioRxiv*. 2022. <https://doi.org/10.1101/2022.09.06.506731>.
- Mulet M, Montaner M, Román D, Gomila M, Kittinger C, Zarfel G, et al. *Pseudomonas* species diversity along the Danube river assessed by rpoD gene sequence and MALDI-TOF MS analyses of cultivated strains. *Front Microbiol*. 2020;11:2114. <https://doi.org/10.3389/fmicb.2020.02114>.
- Nemecek-Marshall M, Laduca R, Fall R. High-level expression of ice nuclei in a *Pseudomonas syringae* strain is induced by nutrient limitation and low temperature. *J Bacteriol*. 1993;175(13):4062–70. <https://doi.org/10.1128/jb.175.13.4062-4070.1993>.
- Nikolić I, Stanković S, Dimkić I, Berić T, Stojšin V, Janse J, et al. Genetic diversity and pathogenicity of *Pseudomonas syringae* pv. *aptata* isolated from sugar beet. *Plant Pathol*. 2018;67(5):1194–207. <https://doi.org/10.1111/ppa.12831>.
- Oueslati M, Mulet M, Zouaoui M, Chandeysson C, Lualcat J, Hajlaoui MR, Berge O, et al. Diversity of pathogenic *Pseudomonas* isolated from citrus in Tunisia. *AMB Express*. 2020;10(1):1–20. <https://doi.org/10.1186/s13568-020-01134-z>.
- Parke JL, Redekar NR, Eberhart JL, Funahashi F. Hazard analysis for *Phytophthora* species in container nurseries: three case studies. *HortTechnology*. 2019;29(6):745–55. <https://doi.org/10.21273/HORTTECH04304-19>.
- Pietsch RB, Vinatzer BA, Schmale DG III. Diversity and abundance of ice nucleating strains of *Pseudomonas syringae* in a freshwater lake in Virginia, USA. *Front Microbiol*. 2017;8:318. <https://doi.org/10.3389/fmicb.2017.00318>.
- Popović T, Ivanović Ž, Trkulja N, Milosavljević A, Ignjatov M. First report of *Pseudomonas syringae* pv. *syringae* on Pea (*Pisum sativum*) in Serbia. *Plant Dis*. 2015a;99(5):724. <https://doi.org/10.1094/PDIS-11-14-1212-PDN>.
- Popović T, Ivanović Ž, Ignjatov M, Milošević D. First report of *Pseudomonas syringae* pv. *coriandricola* causing bacterial leaf spot on carrot, parsley, and parsnip in Serbia. *Plant Dis*. 2015b;99(3):416. <https://doi.org/10.1094/PDIS-10-14-1041-PDN>.
- Popović T, Ivanović Ž, Ignjatov M. First report of *Pseudomonas viridiflava* causing pith necrosis of tomato (*Solanum lycopersicum*) in Serbia. *Plant Dis*. 2015c;99(7):1033. <https://doi.org/10.1094/PDIS-01-15-0052-PDN>.
- Riffaud CH, Morris CE. Detection of *Pseudomonas syringae* pv. *aptata* in irrigation water retention basins by immunofluorescence colony-staining. *Eur J Plant Pathol*. 2002;108(6):539–45. <https://doi.org/10.1023/A:1019919627886>.
- Schneider CA, Rasband WS, Eliceiri KW. NIH image to ImageJ: 25 years of image analysis. *Nat Methods*. 2012;9(7):671–5. <https://doi.org/10.1038/nmeth.2089>.
- Scortichini M, Marcelletti S, Ferrante P, Petriccione M, Firrao G. *Pseudomonas syringae* pv. *actinidiae*: a re-emerging, multi-faceted, pandemic pathogen. *Mol Plant Pathol*. 2012;13(7):631–40. <https://doi.org/10.1111/j.1364-3703.2012.00788.x>.
- Stojšin V, Balaž J, Budakov D, Stanković S, Nikolić I, Ivanović Ž, et al. First report of *Pseudomonas syringae* pv. *aptata* causing bacterial leaf spot on sugar beet in Serbia. *Plant Dis*. 2015;99(2):281. <https://doi.org/10.1094/PDIS-06-14-0628-PDN>.
- Vuković D, Vuković Z, Stanković S. The impact of the Danube Iron Gate Dam on heavy metal storage and sediment flux within the reservoir. *CATENA*. 2014;113:18–23. <https://doi.org/10.1016/j.catena.2013.07.012>.
- Xin XF, Kvitko B, He SY. *Pseudomonas syringae*: what it takes to be a pathogen. *Nat Rev Microbiol*. 2018;16(5):316–28. <https://doi.org/10.1038/nrmicro.2018.17>.
- Yang R, Guan Y, Zhou J, Sun B, Wang Z, Chen H, et al. Phytochemicals from *Camellia nitidissima* chi flowers reduce the Pyocyanin production and motility of *Pseudomonas aeruginosa* PAO1. *Front Microbiol*. 2018;8:2640. <https://doi.org/10.3389/fmicb.2017.02640>.
- Zemunac R, Savic R, Blagojevic B, Benka P, Bezdán A, Salvai A. Assessment of surface and groundwater quality for irrigation purposes in the Danube-Tisa-Danube hydrosystem area (Serbia). *Environ Monit Assess*. 2021;193(8):1–9. <https://doi.org/10.1007/s10661-021-09294-6>.
- Zlatković N, Aleksić G, Gašić K. First report of bacterial canker on blueberry (*Vaccinium corymbosum*) caused by *Pseudomonas syringae* pv. *syringae* in Serbia. *Plant Dis*. 2022;106(3):1057. <https://doi.org/10.1094/PDIS-07-21-1395-PDN>.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions





Genotyping-driven diversity assessment of biocontrol potent *Bacillus* spp. strain collection as a potential method for the development of strain-specific biomarkers

Iva Rosić¹ · Ivan Nikolić¹ · Tamara Ranković¹ · Marina Anteljević¹ · Olja Medić¹ · Tanja Berić¹ · Slaviša Stanković¹

Received: 26 August 2022 / Revised: 1 February 2023 / Accepted: 1 March 2023 / Published online: 13 March 2023
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023

Abstract

Bacillus species are among the most researched and frequently applied biocontrol agents. To estimate their potential as environmentally friendly microbial-based products, reliable and rapid plant colonization monitoring methods are essential. We evaluated repetitive element-based (rep) and Random Amplified Polymorphic DNA (RAPD) PCR (Polymerase Chain Reaction) genotyping in a diversity assessment of 251 strains from bulk soil, straw, and manure samples across Serbia, highlighting their discriminative force and the presence of unique bands. RAPD 272, OPG 5, and (GTG)₅ primers were most potent in revealing the high diversity of a sizable environmental *Bacillus* spp. collection. RAPD 272 also amplified a unique band for a proven biocontrol strain, opening the possibility of Sequence Characterized Amplified Region (SCAR) marker design. That will enable colonization studies using the SCAR marker for its specific detection. This study provides a guide for primer selection for diversity and monitoring studies of environmental *Bacillus* spp. isolates.

Keywords RAPD · Rep · Genotyping · *Bacillus* spp. · Microbial diversity · Biocontrol

Introduction

In light of increasing global awareness of harmful environmental and health effects of agrochemicals, climate change and rapidly growing population, the search for alternative methods of plant growth promotion has never been more needful. Over the years, research on beneficial plant growth-promoting bacteria (PGPB) from the *Bacillus* genus has evinced promising results, especially through their biocontrol mechanism against plant pathogens. Members of the *B. subtilis* complex are of biotechnological importance and several have already been adapted to commercially

available PGPB-based products against bacterial and fungal pathogens (Ferreira et al. 2019).

Crucial to producing and commercializing an environmentally friendly and safe biopesticide is the transition of biocontrol strain efficacy testing from in vitro to in situ testing phase (Marian and Shimizu 2019). Often, some promising biocontrol agents show a lack of in situ activity during the initial phase of testing, usually due to difficulties in establishing and surviving in a new ecological niche (Legein et al. 2020). Therefore, colonization of new habitat is a basic prerequisite for the manifestation of biocontrol activity.

To determine the fate of a potential biocontrol agent in situ, a major challenge is to develop biomarkers for their rapid detection after treatment. Monitoring the population dynamics of the biocontrol agent in the new environment is necessary to conduct efficacy studies and environmental risk assessments and is a prerequisite for the registration of microbial biopesticides (Vilanova et al. 2018; Jiao et al. 2021). In our previous studies, several *Bacillus* strains demonstrated remarkable antimicrobial activity against different phytopathogens (Berić et al. 2012; Dimkić et al. 2013, 2017; Nikolić et al. 2019), highlighting the vast

Communicated by Wen-Jun Li.

Iva Rosić and Ivan Nikolić contributed equally to this work.

✉ Iva Rosić
iva.rosic@bio.bg.ac.rs

¹ University of Belgrade - Faculty of Biology, Center for Biological Control and Plant Growth Promotion, University of Belgrade, Studentski Trg 16, 11000 Belgrade, Serbia

biocontrol potential of *Bacillus* strains and prospective for application as microbial-based products.

Biomarker selection is associated with the identification and determination of *Bacillus* spp. diversity. Over the years, PCR-based technologies have been extensively used to study molecular markers, genetic relationships, and the diversity of *Bacillus* species (Wang et al. 2020). A novel, more informative, and precise genotyping methods, such as Next Generation Multi-Locus Sequence Typing (NG-MLST), often require proficiency in computational analysis (Pérez-Losada et al. 2018). However, genotyping using repetitive sequence-based polymerase chain reaction (rep-PCR), random amplified polymorphic DNA (RAPD-PCR), amplified fragment length polymorphism (AFLP), and pulsed-field gel electrophoresis (PFGE) have over time proved to be important analytical methods that can distinguish between strains of the same bacterial species (Zhang et al. 2021, da Costa Fernandes et al. 2022). RAPD and rep-PCR are equally discriminative alternatives to the bacterial genotyping “gold standard” PFGE (Stefańska et al. 2022). In addition, they have the advantages of being easy to perform, fast, and cost-effective methods (Sharma et al. 2020). Rep-PCR genotyping generates multiple amplicons of intergenic DNA between binding sites of adjacent repetitive elements. Binding sites for rep-PCR often include (GTG)₅ binding site, enterobacterial repetitive intergenic consensus sequences (ERIC), repetitive extragenomic palindromic sequences (REP) and BOX elements (Rodríguez et al. 2019; Macůrková et al. 2021). On the other hand, RAPD-PCR employs single arbitrary oligonucleotide primers that generate random amplicons; thus, prior sequence knowledge is unnecessary. Although RAPD-PCR was developed three decades ago (Williams et al. 1990) and did not involve sequencing, it is still widely used for genotyping due to its versatility and robustness. RAPD-PCR has also been extensively described as a successful tool for designing strain-specific molecular markers for environmental *Bacillus* spp. isolates (Reddypriya et al. 2019; Hernández et al. 2020; Orce et al. 2021). In addition, those methods have important applications in the design of RAPD/rep-derived molecular strain-specific or species-specific markers, for example, SCAR (Sequence Characterized Amplified Region) primers (Bhagyawant 2016; Di Francesco et al. 2018). Conversion of unique RAPD/rep amplicons into SCAR primers enables specific detection and quantification of certain strains and overcomes the limitation of RAPD genotyping concerning reproducibility (Huang et al. 2018; Hu et al. 2021). The SCAR markers could serve as a prominent tool for monitoring the colonization capabilities of targeted biocontrol agents. Therefore, genotyping-based diversity determination could potentially develop powerful tools to rapidly and reliably detect and evaluate the effectiveness of

plant colonization by PGPB (Ambreetha and Balachandar 2023).

This study aimed to assess genotyping-based diversity and compare discriminative forces of rep-PCR and RAPD-PCR genotyping of 251 *Bacillus* spp. strains, isolated from various natural samples across Serbia. Therefore, determining a fingerprinting approach with the finest discriminative force could be important in future studies regarding strain-specific marker design. Consequently, they serve as a suitable basis for developing a valid platform to evaluate the colonization capabilities among *Bacillus* spp. biocontrol strains.

Materials and methods

Bacterial strains and culture conditions

The *Bacillus* spp. isolates used in this study are part of a collection belonging to the Laboratory of Microbiology, Faculty of Biology. From different localities across Serbia, 153 strains were isolated from bulk soil, 34 strains from straw, and 64 strains from manure (Berić et al. 2009; Stanković et al. 2012). The isolation method was based on the ability of *Bacillus* spp. to form resistant endospores under unfavorable physical and chemical conditions. The vegetative cells in the samples were thermally inactivated by heating 1 g of the sample in 1 ml Nutrient broth (NB) at 80 °C for 10 min. Samples were then plated on Luria–Bertani agar (LA) and incubated at 30 °C for 48 h until pure cultures were obtained for each isolate. Bacterial isolates were characterised at the genus level by microscopic appearance, Gram staining and catalase test (Berić et al. 2012). Isolates were revitalized in Luria–Bertani (LB) broth at 30 °C for 24 h on a rotary shaker at 180 rpm. Overnight cultures were plated on LA medium and incubated at room temperature for 48 h until pure colonies were obtained for each strain. Pure cultures were stored in 25% glycerol and kept at – 80 °C.

Genomic DNA extraction and quantification

For DNA isolation, strains were incubated on LA plates at 30 °C for 24 h. DNA extraction was performed using the CTAB-chloroform method based on Le Marrec et al. (2000) protocol, adapted for Gram-positive bacteria. Colonies were suspended in lysis buffer [TE buffer, pH 7.6 (10 mM TRIS; 1 mM EDTA; dH₂O) with the addition of lysozyme (20 mg ml⁻¹)] and incubated at 37 °C for 30 min. Then, 100 µl 5 M NaCl and 300 µl 3% CTAB buffer, pH 8.0 (CTAB; 1 M Tris; 1.4 M NaCl; 0.5 M EDTA; dH₂O; PVP) were added to lysate. After incubation at 65 °C for 20 min, 800 µl of chloroform was added and the solution was

centrifuged at 13,000 rpm for 15 min. In separate tubes, 1/10 of the total supernatant volume of Na-acetate (pH 5) and an equal volume of ice-cold isopropyl alcohol were mixed with the supernatant. After centrifugation (13,000 rpm, 15 min) pellet was left to dry in a heating block at 37 °C for 30 min. Pellet was then resuspended in a 50 µl TE buffer with the addition of RNase [TE buffer 49 µl; RNase (10 mg ml⁻¹) 1 µl]. Samples were incubated for 15 min at 37 °C. DNA concentration was measured spectrophotometrically (NanoDrop 2000c, Thermo Fisher Scientific, USA) and 50 ng µl⁻¹ DNA was prepared. Samples were stored at - 20 °C before use.

PCR conditions

The primers sequences (Table 1) and modified thermal cycling conditions used in *Bacillus* spp. genotyping were presented in Table 2. Genotyping using BOX and (GTG)₅ primers were performed in a 25 µl reaction mixture containing: 10 µl OneTaq Quick-Load 2×Master Mix with

Standard Buffer (NEB, USA); 1.5 µl BOX A1R primer; 7.5 µl distilled DNase/RNase free PCR water (Gibco, ThermoFisher Scientific, USA) and 1 µl genomic DNA in the concentration of 50 ng µl⁻¹. ERIC amplification was performed in 25 µl reaction mix containing: 2.5 µl 10×KAPA Taq buffer (KAPA Biosystems Inc, USA); 2.5 µl MgCl₂ (25 mM); 0.5 µl dNTP mix (2.5 mM per nucleotide); 1.9 µl of each primer (10 mM); 15.6 µl PCR water; 0.1 µl KAPA Taq Polymerase (5 U µl⁻¹, KAPA Biosystems Inc, USA) and 1 µl gDNA.

The amplification with RAPD 272 primer was performed in 25 µl reaction mix containing: 2.5 µl 10×KAPA Taq buffer; 2.3 µl MgCl₂; 0.63 µl dNTP mix; 1.9 µl of each primer; 16.6 µl PCR water; 0.1 µl KAPA Taq Polymerase and 1 µl gDNA. DNA amplification with OPB07, OPO02 and OPG 5 primers was performed in 25 µl reaction mixture containing: 2.5 µl 10×KAPA Taq buffer; 1.5 µl MgCl₂; 2 µl dNTP mix; 2 µl primers; 15.85 µl PCR water; 0.15 µl KAPA Taq Polymerase and 1 µl gDNA. The 20 µl reaction mixture for RAPD 1247 consisted of: 10 µl OneTaq Quick-Load

Table 1 Sequences of primers used in this study

Primers	Sequence	References
BOX A1R	CTACGGCAAGGCGACGCTGACG	Allipi and Aguilar (1998)
ERIC 1R	ATGTAAGCTCCTGGGGAT	Freitas et al. (2008)
ERIC 2	AAGTAAGTGACTGGGGTG	Freitas et al. (2008)
(GTG) ₅	GTGGTGGTGGTGGTG	Versalovic et al. (1998)
RAPD 272	AGCGGGCCAA	Hematzadeh and Haghkhal (2021)
OPB 07	GGTGACGCAG	Khawal et al. (2017)
OPO 02	ACGTAGCGTC	Khawal et al. (2017)
OPG 5	CTGAGACGGA	Felici et al. (2008)
RAPD 1247	AAGAGCCCGT	Gallori et al. (1998)

Table 2 Modified rep-PCR and RAPD-PCR protocols

Primers	PCR conditions				
	Initial denaturation	Number of cycles			Final elongation
		Denaturation	Hybridization	Elongation	
BOX A1R; ERIC1R; ERIC 2	95 °C 7 min	× 30			65 °C 16 min
		94 °C 1 min	52 °C 1 min	65 °C 8 min	
(GTG) ₅	94 °C 5 min	× 35			72 °C 1 min
		94 °C, 30 s	45 °C, 1 min	72 °C, 1 min	
RAPD 272	94 °C 2 min	× 35			72 °C 10 min
		94 °C, 30 s	35 °C, 30 s	72 °C, 2 min	
OPB 07; OPO 02	94 °C 1 min	× 45			72 °C 3 min
		94 °C, 1 min	35 °C, 30 s	72 °C, 1 min	
OPG 5	94 °C 4 min	× 30			72 °C 5 min
		94 °C, 1 min	36 °C, 1 min	72 °C, 30 s	
RAPD 1247	94 °C 1 min	× 30			72 °C 10 min
		94 °C, 1 min	36 °C, 1 min	72 °C, 2 min	

2×Master Mix with Standard Buffer; 2 µl primer and 7 µl PCR water. All amplification reactions were performed with MiniAmp™ Thermal Cycler (Thermo Fisher Scientific, USA).

Gel electrophoresis and visualization of genetic profiles

PCR products were mixed with loading dye (0.25 g bromophenol blue; 6 ml 50% glycerol; 4 ml dH₂O) and visualized on 1% agarose gel containing an aqueous solution of 10 mg ml⁻¹ ethidium-bromide (SERVA, Germany). Electrophoresis was run in 0.5×Tris–borate-EDTA buffer (5×TBE: 54 g Tris; 27.5 g Boric acid; 40 ml 0.5 M pH 8 EDTA; dH₂O to 1 L) at a constant voltage of 90 V and 300 mA amperage for 90 min.

The gels were visualized using a transilluminator (LKB, Transilluminator 2011 Macrovue UV Light, Sweden).

Data analysis

Obtained genetic profiles were first compared using GelAnalyzer 19.1 software. Profiles were then converted into a binary matrix where digit “1” signifies the presence of a band on a certain position and “0” bands absence. Data were analyzed using a formula for polymorphism calculation, modified and adapted from Bardakci and Skibinski (1994):

$$P_{xy} = n_{xy} / (n_x + n_y).$$

Variable n_{xy} presents the number of polymorphic bands between x and y strain, while n_x and n_y are the total numbers of bands per strain. Polymorphism generated with each primer was calculated as the average of P_{xy} between all compared strains. To evaluate typing efficiency of the eight different primers, a discrimination index (D) was calculated as described by Hunter and Gaston (1988). Cluster analysis was performed in CLIQS 1D software (TotalLab, UK), applying the unweighted pair group method with arithmetic mean (UPGMA). Relations between different profiles were presented as dendrograms.

As means of comparing the convenience of each typing method, we considered the following parameters: (I) Total number of bands; (II) Size range of bands; (III) Number of different profiles; (IV) Polymorphism index; (V) Discrimination index (D index) and (VI) Number of unique bands generated with each primer.

Results and discussion

The discriminative abilities of 8 different rep-PCR and RAPD primers were compared to estimate the diversity of 251 *Bacillus* spp. strains. Comparing rep-PCR binding patterns revealed 48 different genetic profiles generated by BOX A1R primer, 55 with (GTG)₅ and 30 with ERIC 1R and ERIC 2 primers. Grouping of *Bacillus* spp. strains into profiles after the amplification reaction with RAPD primers was as follows: 34 genetic profiles generated by RAPD 1247 and OPB07, 44 by RAPD 272, 19 by OPO02 and 55 by OPG 5 (Tables S1–S8; Fig. S1–S8).

According to the assigned cut-off value of 0.2, similar profiles were grouped in clusters, representing strains relatedness. The highest number of separate clusters was achieved by RAPD 272 primer (Fig. 1A) and the least with ERIC and RAPD 1247 (Fig. 1B).

It was further observed that amplification products with BOX A1R were absent in 19 strains, 70 with ERIC primer and only one with (GTG)₅. The highest number of strains without amplicons was perceived in reaction with RAPD 1247 (more than half of the strains). At the same time, amplification using OPG 5 failed to produce bands in only nine strains. OPO02, OPB07 and RAPD 272 did not yield amplification for 67, 65 and 17 strains, respectively.

Polymorphism was calculated separately for three environmental sources of isolates and then generally for the whole collection as a mean value. The level of discrimination or probability that any two randomly chosen strains will be of the same type was calculated. Based on the final results summarized in Table 3, it was observed that RAPD 272 primer demonstrated the highest discriminative force in *Bacillus* spp. genotyping, whereas RAPD 1247 scored the lowest according to the selected parameters. The D index for all methods was higher than 0.9. Therefore, typing results can be interpreted with confidence (Hunter and Gaston 1988). Accordingly, we perceived RAPD 272 genotyping method as the most successful in revealing the differences among *Bacillus* spp. isolates. Even though OPG 5 showed very similar results to RAPD 272, the number of bands, size, and ability to distinguish between similar strains favored RAPD 272.

Conversely, RAPD 1247 primer was the least effective in *Bacillus* spp. genotyping as more than 50% of the reactions were negative. The PCR program was optimized and tested, and all the negative reactions were repeated, leading to the same result. Therefore, we are convinced that technical issues were less likely to cause amplification absence in the strains. RAPD 1247 genotyping of *Bacillus* spp. was successfully applied only in Batinić et al. (1997) and Gallori et al. (1998), on a much smaller sample size. In addition, novel data employing this particular method on *Bacillus* spp.




Fig. 1 *Bacillus* spp. collection genotyping and diversity estimation. **A** Dendrogram showing relatedness of grouped strains (profiles) where similar profiles were divided into 10 different clusters after RAPD

272-PCR. **B** Dendrogram showing relatedness of grouped strains (profiles) where similar profiles were divided into 5 different clusters after RAPD 1247-PCR

is lacking. Consequently, the obtained results could not be compared to form an accurate conclusion.

ERIC-PCR, a widely used genotyping method for *Escherichia coli* and *Salmonella typhimurium*, for which ERIC elements were first described (Versalovic et al. 1991), was also less effective. Most of the studies on *Bacillus* spp. were done on *B. cereus* group species and results varied. Some reported a higher discriminative force of ERIC compared to other rep-PCR methods (Subbanna et al. 2018), while in a greater number of studies, BOX and (GTG)₅ proved to be more effective (Getahun et al. 2020). Avsar et al. (2017) and Orce et al. (2021) reported that a higher degree of genetic diversity was revealed by RAPD genotyping compared to ERIC, which also supports our finding. Furthermore, RAPD was described as the best method for molecular typing of *Bacillus* spp. (Gupta and

Joia 2016). However, even though RAPD 272 primer is extensively used in other genera (*Pseudomonas* spp. typing, for example), it has been rarely applied to *Bacillus* spp. (Emami et al. 2019; Hematizadeh and Haghkhal 2021).

In addition, we compared each primer's ability to assess genetic diversity for the natural sources of isolates separately (Fig. 2). For straw isolates genotyping, OPO 02, followed by RAPD 272 and BOX, according to polymorphism value, could be the primers of choice in diversity assessment. As for soil and manure isolates, RAPD 272, (GTG)₅ and OPG 5 -PCR were the best methods. Considering the unequal number of isolates from each natural sample and the number of negative reactions of the same strains, reported results should be taken with a reserve and need further exploration.


A variety of experiments were conducted with this collection of *Bacillus* spp. The results underlined the

Table 3 Summary of all data obtained with eight different primers

	Total No. of bands	Size of bands (kb)	Polymorphism (mean value)	D index	No. of unique bands	Strain code and unique band size (kb)
RAPD 272	1 - 8	0.5 - 9	0.7171	0.9683	1	SS- 12.6.1/12.6.2 (0.5)
(GTG) ₅	1 - 7	0.4 - 10	0.7186	0.9478	2	SS- 32.6 (0.5); SS- 9.3 (0.6)
OPG5	1 - 6	0.1 - 3	0.7152	0.9676	2	SS- 38.4.1 (3); SS- 6.6.1/6.4 (0.3)
BOX	1 - 9	0.1 - 10	0.6925	0.9619	0	None
OPO 02	1 - 5	0.7 - 10	0.6032	0.9253	1	SS- 8.1.2 (0.5)
ERIC	1 - 6	0.2 - 3	0.5185	0.9470	0	None
OPB 07	1 - 6	0.2 - 3	0.4829	0.9390	0	None
1247	1 - 5	0.4 - 3	0.4332	0.9392	0	None

Red-colored cells indicate the highest value, while green is the lowest. Strains SS-12.6.1 and SS-12.6.2, as well as SS-6.6.1 and SS-6.4, were considered clones due to being isolated from the same location and having the same genetic profiles

Fig. 2 Comparison of primers' force in estimating genetic diversity of *Bacillus* spp. isolates from straw (green), soil (yellow), and manure (blue). OPO 02 primer showed the highest potential in the differentiation of straw isolates, whereas (GTG)₅ in both soil and manure isolates. RAPD 272 demonstrated equally good discriminative power for all three sources (colour figure online)



plant growth-promoting features of several strains and their immense potential as BCAs. Berić et al. (2012) study revealed that concentrated cell-free culture supernatants of 127 *Bacillus* strains exhibited antagonistic activity against five plant pathogens: *Burkholderia glumae*, *Burkholderia cepacia*, *Erwinia carotovora*, *Pseudomonas fuscovaginae* and *Xanthomonas oryzae*. Moreover, a strong biocontrol activity of lipopeptide extracts from the five *Bacillus* strains against two pathovars *P. syringae* pv. *aptata* and *Xanthomonas arboricola* pv. *juglandis* was indicated in Dimkić et al. (2013) and Dimkić et al. (2017). In vivo application of one strain (*B. amyloliquefaciens* SS-12.6) showed an antifungal effect, as well. Nikolić et al. (2019) successfully demonstrated the biocontrol activity of crude lipopeptide extracts of *B. amyloliquefaciens* (SS-12.6 and SS-38.4.1) and *B. pumilus* (SS-10.7) strains against *P. syringae* pv. *aptata* in vitro and in planta.

Both rep and RAPD methods produced strain-specific profiles suggesting a great potential for designing strain-specific biomarkers (Table 3). The four primers produced six unique bands (Fig. S1-S4), a notable result considering the number of strains in this study. No more than 20 strains were commonly compared to identify BCAs by SCAR markers successfully (Gotor-Vila et al. 2016). We also observed many strain-specific profiles; however, the positions of individual bands overlapped with those of other profiles. The most significant result was the detection of species-specific profiles for proven biocontrol strains. For *B. amyloliquefaciens* SS- 12.6 (Fig. S1) with RAPD 272 and *B. amyloliquefaciens* SS- 38.4.1 with OPG 5 primer (Fig. S3). The unique band of 0.5 kb obtained for SS- 12.6.1/SS-12.6.2 could be used in SCAR marker design for specific detection and quantification of biocontrol strain after application in planta. That is

necessary for estimating an optimal number of bacteria that should be maintained for biocontrol activity effects to manifest. Obtained results could be essential in developing strain-based commercial products as guidance for their application (Ku et al. 2021). Strain-specific band for SS-38.4.1 (size 3 kb) could be less likely the choice as most of the rep or RAPD PCR products for SCAR markers design are no longer than 0,8 kb (Felici et al. 2008). In this regard (GTG)₅ could be considered the most promising for strain-specific primer development, with two unique bands of optimal size.

Primers OPO 02 and OPG 5 have already been proven to generate polymorphic bands for SCAR marker design (Felici et al. 2008; Khowal et al. 2017). To the best of our knowledge, primers RAPD 272 and (GTG)₅ have not yet been used for SCAR marker design in *Bacillus* spp. On top of that, the unique band obtained with RAPD 272 primer belongs to SS-12.6.1/12.6.2 strains characterized as *B. amyloliquefaciens*, which demonstrated antimicrobial and antifungal activity against plant pathogens (Dimkić et al. 2013). Those results could be of great importance for further studies on the possible development of strain-based biocontrol products. The necessary step towards that goal would be a design of RAPD 272-based SCAR primers and their SS-12.6.1/12.6.2 specificity validation throughout the whole collection. Specific identification of biocontrol strains would enable its precise *in planta* detection among the abundance of indigenous microbial species (Zhang et al. 2020). Strain monitoring using quantitative or digital PCR techniques in colonization quantification assays would evaluate the commercial potential of SS-12.6.1/12.6.2 biocontrol agent.

Conclusion

Rep and RAPD PCR are still the epitome of both rapid and economical methods for diversity assessments of large strain collections. Based on the comparison of various factors, two RAPD primers, RAPD 272 and OPG 5 and one rep-primer (GTG)₅ can be considered as primers of choice in the genetic diversity assessment of natural *Bacillus* spp. isolates. In addition, this study provides a starting point for the design of SCAR markers for biocontrol strains in the collection based on unique bands obtained with RAPD and rep primers. Future studies will include cloning and sequencing unique amplicons and validating the specificity of the designed SCAR markers. Selected SCAR markers will be used in biocontrol experiments *in planta* for quantification of the colonization potential of biocontrol strains.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00203-023-03460-9>.

Author contributions All authors contributed to the study's conception and design. Conceptualization and methodology were designed by IN and OM. IR, MA, and TR: performed the experiments, and collected, analyzed, and visualized data. The original draft was written by IR and IN. TB and SS: reviewed and edited the draft. All authors read and approved the final manuscript.

Funding This work was supported by the Ministry of Science, Technological Development and Innovations of the Republic of Serbia, grant numbers 451-03-47/2023-01/200178 and Center for leadership development, Belgrade, Republic of Serbia, grant number: 2020-5. Authors Ivan Nikolić, Olja Medić and Tamara Ranković received grant 2020-5, while all authors received grant 451-03-47/2023-01/200178.

Data Availability The datasets generated and analysed during this study are available from the corresponding author on the request.

Declarations

Conflict of interest The authors declare no conflict of interest in the publication of this manuscript.

References

- Alippi AM, Aguilar OM (1998) Characterization of isolates of *Paenibacillus larvae* subsp. *larvae* from diverse geographical origin by the polymerase chain reaction and BOX primers. *J Invertebr Pathol* 72:21–27. <https://doi.org/10.1006/jipa.1998.4748>
- Ambreetha S, Balachandar D (2023) SCAR marker: a potential tool for authentication of agriculturally important microorganisms. *J Basic Microbiol* 63:4–16. <https://doi.org/10.1002/jobm.202200419>
- Avşar C, Koyuncu H, Aras ES (2017) Isolation and molecular characterization of *Bacillus* spp. isolated from soil for production of industrial enzymes. *Biol Chem Res* 3:72–86
- Bardakci F, Skibinski DOF (1994) Application of the RAPD technique in tilapia fish: species and subspecies identification. *Heredity* 73:117–123. <https://doi.org/10.1038/hdy.1994.110>
- Batinić T, Schmitt J, Ulrike M, Schulz D (1997) Construction of RAPD-generated DNA probes for the quantification of *Bacillus subtilis* FZB C and the evaluation of its biocontrol efficiency in the system *Cucumis sativus* - *Pythium ultimum*. *J Plant Dis Prot* 105:168–180
- Berić T, Urdaci MC, Stanković S, Knežević-Vukčević J (2009) RAPD analysis of genetic diversity and qualitative assessment of hydrolytic activities in a collection of *Bacillus* sp. isolate. *Arch Biol Sci* 61:645–652. <https://doi.org/10.2298/ABS0904645B>
- Berić T, Kojić M, Stanković S, Topisirović L, Degrossi G, Myers M, Venturi V, Fira Đ (2012) Antimicrobial activity of *Bacillus* sp. natural isolates and their potential use in the biocontrol of phytopathogenic bacteria. *Food Technol Biotechnol* 50:25–31
- Bhagyawant SS (2016) RAPD-SCAR markers: an interface tool for authentication of traits. *Int J Biosci Med* 4:1–9. <https://doi.org/10.4236/jbm.2016.41001>
- Da Costa FG, de Prado Costa DK, de Oliveira NS, de Sousa ECP, Machado DHB, Polanczyk RA, de Siqueira HAA, da Silva MC (2022) Genetic diversity of Brazilian *Bacillus thuringiensis* isolates with toxicity against *Aedes aegypti* (Diptera: Culicidae). *Sci Rep* 12:1–9. <https://doi.org/10.1038/s41598-022-18559-0>

- Di Francesco A, Calassanzio M, Ratti C, Mari M, Folchi A, Baraldi E (2018) Molecular characterization of the two postharvest biological control agents *Aureobasidium pullulans* L1 and L8. *Biol Control* 123:53–59. <https://doi.org/10.1016/j.biocontrol.2018.05.005>
- Dimkić I, Živković S, Berić T, Ivanović Ž, Gavrilović V, Stanković S, Fira Đ (2013) Characterization and evaluation of two *Bacillus* strains, SS-12.6 and SS-13.1, as potential agents for the control of phytopathogenic bacteria and fungi. *Biol Control* 65:312–321. <https://doi.org/10.1016/j.biocontrol.2013.03.012>
- Dimkić I, Stanković S, Nišavić M, Petković M, Ristivojević P, Fira Đ, Berić T (2017) The profile and antimicrobial activity of *Bacillus* lipopeptide extracts of five potential biocontrol strains. *Front Microbiol* 8:1–12. <https://doi.org/10.3389/fmicb.2017.00925>
- Emami A, Keshavarzi A, Pirbonyeh N, Behbahani MR (2019) Identification of different faces of *Pseudomonas aeruginosa* isolates in burn patients by genetic fingerprinting. *Gene Reports* 15:100377. <https://doi.org/10.1016/j.genrep.2019.100377>
- Felici C, Vettori L, Toffanin A, Nuti M (2008) Development of a strain-specific genomic marker for monitoring a *Bacillus subtilis* biocontrol strain in the rhizosphere of tomato. *FEMS Microbiol Ecol* 65:289–298. <https://doi.org/10.1111/j.1574-6941.2008.00489.x>
- Ferreira CMH, Soares HMVM, Soares EV (2019) Promising bacterial genera for agricultural practices: an insight on plant growth-promoting properties and microbial safety aspects. *Sci Total Environ* 682:779–799. <https://doi.org/10.1016/j.scitotenv.2019.04.225>
- Freitas DB, Reis MP, Lima-Bittencourt CI, Costa PS, Assis PS, Chartone-Souza E, Nascimento AM (2008) Genotypic and phenotypic diversity of *Bacillus* spp. isolated from steel plant waste. *BMC Res Notes* 1:1–11. <https://doi.org/10.1186/1756-0500-1-92>
- Gallori E, Franchi M, Rinaldi L, Vettori C (1998) Interspecific transformation of *Bacillus subtilis* by clay-bound DNA in non-sterile soil. *Symbiosis* 25:311–322
- Getahun A, Kiros S, Muleta D, Assefa F (2020) Genetic and metabolic diversities of rhizobacteria isolated from degraded soil of Ethiopia. *Heliyon* 6:e05697. <https://doi.org/10.1016/j.heliyon.2020.e05697>
- Gotor-Vila A, Teixidó N, Usall J, Dashevskaya S, Torres R (2016) Development of a SCAR marker and a strain-specific genomic marker for the detection of the biocontrol agent strain CPA-8 *Bacillus amyloliquefaciens* (formerly *B. subtilis*). *Ann Appl Biol* 169:248–256. <https://doi.org/10.1111/aab.12298>
- Gupta A, Joia J (2016) Microbes as potential tool for remediation of heavy metals: a review. *J Microbial Biochem Technol* 8:364–372. <https://doi.org/10.4172/1948-5948.1000310>
- Hematzadeh A, Haghkhal M (2021) Biotyping of isolates of *Pseudomonas aeruginosa* isolated from human infections by RAPD and ERIC-PCR. *Heliyon* 7:e07967. <https://doi.org/10.1016/j.heliyon.2021.e07967>
- Hernández I, Sant C, Martínez R, Fernández C (2020) Design of bacterial strain-specific qPCR assays using NGS data and publicly available resources and its application to track biocontrol strains. *Front Microbiol* 11:1–11. <https://doi.org/10.3389/fmicb.2020.00208>
- Hu J, Yao Y, Yang Y, Gao C, Zhang F, Xia R, Ran C, Zhang Z, Liu CJ, Zhou Z (2021) Advances of intra-species molecular typing analysis of aquatic probiotics approved by the Chinese Ministry of Agriculture. *Rev Aquat Sci* 13:178–188. <https://doi.org/10.1111/raq.12469>
- Huang CH, Liou JS, Huang L, Watanabe K (2018) Developing novel species-specific DNA markers for PCR-based species identification of the *Lactobacillus sakei* group. *Lett Appl Microbiol* 66:138–144. <https://doi.org/10.1111/lam.12825>
- Hunter PR, Gaston MA (1988) Numerical index of the discriminatory ability of typing systems: an application of Simpson's index of diversity. *J Clin Microbiol* 26:2465–2466. <https://doi.org/10.1128/jcm.26.11.2465-2466.1988>
- Jiao X, Takishita Y, Zhou G, Smith DL (2021) Plant associated rhizobacteria for biocontrol and plant growth enhancement. *Front Plant Sci* 12:634796. <https://doi.org/10.3389/fpls.2021.634796>
- Khowal S, Siddiqui MZ, Ali S, Khan MT, Khan MA, Naqvi SH, Wajid S (2017) A report on extensive lateral genetic reciprocation between arsenic resistant *Bacillus subtilis* and *Bacillus pumilus* strains analyzed using RAPD-PCR. *Mol Phylogenet Evol* 107:443–454. <https://doi.org/10.1016/j.ympev.2016.12.010>
- Ku Y, Yang N, Pu P, Mei X, Cao L, Yang X, Cao C (2021) Biocontrol mechanism of *Bacillus subtilis* C3 against bulb rot disease in *Fritillaria taipaiensis* P.Y.Li. *Front Microbiol* 12:1–14. <https://doi.org/10.3389/fmicb.2021.756329>
- Le Marrec C, Hyronimus B, Bressollier P, Verneuil B, Urdaci MC (2000) Biochemical and genetic characterization of coagulin a new antilisterial bacteriocin in the pediocin family of bacteriocins produced by *Bacillus coagulans* I4. *Appl Environ Microbiol* 66:5213–5220. <https://doi.org/10.1128/AEM.66.12.5213-5220.2000>
- Legein M, Smets W, Vandenneuvel D, Eilers T, Muysshondt B, Prinsen E, Samson R, Lebeer S (2020) Modes of action of microbial biocontrol in the phyllosphere. *Front Microbiol* 11:1619. <https://doi.org/10.3389/fmicb.2020.01619>
- Macůrková A, Maryška L, Jindřichová B, Drobníková T, Vrchotová B, Pospíchalová R, Záruba K, Hubáček T, Siegel J, Burketová L, Lovecká P, Valentová O (2021) Effect of round-shaped silver nanoparticles on the genetic and functional diversity of soil microbial community in soil and “soil-plant” systems. *Appl Soil Ecol* 168:104165. <https://doi.org/10.1016/j.apsoil.2021.104165>
- Marian M, Shimizu M (2019) Improving performance of microbial biocontrol agents against plant diseases. *J Gen Plant Pathol* 85:329–336. <https://doi.org/10.1007/s10327-019-00866-6>
- Nikolić I, Berić T, Dimkić I, Popović T, Lozo J, Fira Đ, Stanković S (2019) Biological control of *Pseudomonas syringae* pv. *aptata* on sugar beet with *Bacillus pumilus* SS-10.7 and *Bacillus amyloliquefaciens* (SS-12.6 and SS-38.4) strains. *J Appl Microbiol* 126:165–176. <https://doi.org/10.1111/jam.14070>
- Orce IG, Martínez FL, Aparicio M, Torres MJ, Rajal VB, Irazusta VP (2021) Genetic fingerprint and diversity evaluation of halophilic *Bacillus* species by RAPD-PCR. *An Acad Bras Cienc* 93:1–8. <https://doi.org/10.1590/0001-376520210191430>
- Pérez-Losada M, Arenas M, Castro-Nallar E (2018) Microbial sequence typing in the genomic era. *Infect Genet Evol* 63:346–359. <https://doi.org/10.1016/j.meegid.2017.09.022>
- Reddy Priya P, Soumare A, Balachandar D (2019) Multiplex and quantitative PCR targeting SCAR markers for strain-level detection and quantification of biofertilizers. *J General Microbiol* 59:111–119. <https://doi.org/10.1002/jobm.201800318>
- Ruiz Rodríguez LG, Mohamed F, Bleckwedel J, Medina R, De Vuyst L, Hebert EM, Mozzi F (2019) Diversity and functional properties of lactic acid bacteria isolated from wild fruits and flowers present in northern Argentina. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2019.01091>
- Sharma A, Lee S, Park YS (2020) Molecular typing tools for identifying and characterizing lactic acid bacteria: a review. *Food Sci Biotechnol* 29:1301–1318. <https://doi.org/10.1007/s10068-020-00802-x>
- Stanković S, Mihajlović S, Draganić V, Dimkić I, Vukotić G, Berić T, Fira Đ (2012) Screening for the presence of biosynthetic genes for antimicrobial lipopeptides in natural isolates of *Bacillus* sp. *Arch Biol Sci* 64:1425–1432. <https://doi.org/10.2298/ABS1204425S>
- Stefańska I, Kwiecień E, Górzynska M, Sałamaszyńska-Guz A, Rzewuska M (2022) RAPD-PCR-Based fingerprinting method

- as a tool for epidemiological analysis of *Trueperella pyogenes* infections. *Pathogens* 11:562. <https://doi.org/10.3390/pathogens11050562>
- Subbanna ARNS, Khan MS, Srivastava RM, Mishra PK, Babu BK, Venkateswarlu V (2018) Interspecies diversity of *Bacillus thuringiensis* isolates native from North Western Indian Himalayas. *J Environ Biol* 39:306–313. <https://doi.org/10.22438/jeb/39/3/MRN-574>
- Versalovic J, De Bruijn FJ, Lupski JR (1998) Repetitive sequence-based PCR (rep-PCR) DNA fingerprinting of bacterial genomes. *Bact Genomes*. https://doi.org/10.1007/978-1-4615-6369-3_34
- Versalovic J, Koeuth T, Lupski JR (1991) Distribution of repetitive DNA sequences in eubacteria and application to finerprinting of bacterial genomes. *Nucleic Acids Res* 19:6823–6831. <https://doi.org/10.1093/nar/19.24.6823>
- Vilanova L, Teixidó N, Usall J, Balsells-Llauradó M, Gotor-Vila A, Torres R (2018) Environmental fate and behaviour of the biocontrol agent *Bacillus amyloliquefaciens* CPA-8 after preharvest application to stone fruit. *Pest Manag Sci* 74:375–383. <https://doi.org/10.1002/ps.4716>
- Wang Z, Yu ZX, Solanki MK, Yang LT, Xing YX, Dong DF, Li YR (2020) Diversity of sugarcane root-associated endophytic *Bacillus* and their activities in enhancing plant growth. *J Appl Microbiol* 128:814–827. <https://doi.org/10.1111/jam.14512>
- Williams JGK, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res* 18:6531–6535. <https://doi.org/10.1093/nar/18.22.6531>
- Zhang S, Ma Y, Jiang W, Meng L, Cao X, Hu J, Chen J, Li J (2020) Development of a strain - specific quantification method for monitoring *Bacillus amyloliquefaciens* TF28 in the rhizospheric soil of soybean. *Mol Biotechnol* 62:521–533. <https://doi.org/10.1007/s12033-020-00268-6>
- Zhang C, Yu X, Wang D, Gui Y, Wang C, Li Q, Wang J, Yin B, Pan Z, Gu R (2021) Rapid strain-specific identification of two *Lactobacillus rhamnosus* strains using PCR based on gene family analysis. *Lwt* 146:111395. <https://doi.org/10.1016/j.lwt.2021.111395>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com

GAPP24

The presence of ice nucleation active *Pseudomonas syringae* in the Danube River basin

Marina Anteljević^{1*}, Iva Rosić¹, Tamara Ranković¹, Olja Medić¹, Tanja Berić¹, Slaviša Stanković¹ and Ivan Nikolić¹

¹ University of Belgrade, Faculty of Biology, Center for Biological Control and Plant Growth Promotion, Studentski trg 16, Belgrade, Serbia

* Correspondence: marina.anteljevic@bio.bg.ac.rs

Pseudomonas syringae (*Psy*) is a widespread complex of plant pathogenic bacteria that can provide nutrients by causing frost damage to plant tissue. They synthesize the ice nucleation protein InaZ, which is anchored to the outer membrane of the bacterial cell. It enables ice formation by bringing water molecules into an "ice-like" arrangement at temperatures as high as -2°. The aim of our study was to investigate the ability of *Psy* isolates to nucleate ice. The ice nucleation activity (INA) test was performed on 51 *Psy* representatives from the Danube River Basin in Serbia, which is an important irrigation source. The isolates were exposed to decreasing temperatures from -2° to -10° in a decline of 1°. The INA tests revealed 39 active strains (76.5% of the collection). Depending on the ice-forming temperature, three different INA types were observed: warmer than -4° (type I), -4° to -7° (type II), and colder than -7° (type III). Based on the average freezing temperatures, seven isolates belonged to type I, 14 to type II and 18 to type III. The highest temperature at which ice formed was -3°C. Our study showed the presence of INA bacteria in the irrigation water source, which may increase the risk of frost damage to plants. Frost damage causes significant losses to frost-sensitive plants, so its prevention is essential for crop health.

Keywords: *Pseudomonas syringae*; plant pathogen; ice nucleation activity; frost damage.

GAPP25

Screening of AHL lactonase activity in *Bacillus* spp. strains isolated from different natural samples

Iva Rosić¹, Ivan Nikolić¹, Tamara Ranković¹, Marina Anteljević¹, Tanja Berić¹, Slaviša Stanković¹ and Olja Medić¹

¹ University of Belgrade, Faculty of Biology, Center for Biological Control and Plant Growth Promotion, 11000 Belgrade, Serbia

* Correspondence: iva.rosic@bio.bg.ac.rs

Through evolution, pathogenic bacteria developed mechanism to increase their virulence to hosts by production of *quorum sensing* (QS) signaling molecules, primarily acyl homoserine-lactones (AHLs). Since the discovery, QS phenomenon has been promising target in biocontrol strategies involving *quorum quenching* (QQ) mechanisms. The great potential in this area lies in detection, characterization and possible application of AHL-lactonases from *Bacillus* spp. The purpose of this study was to test presence and activity of AHL-lactonases in natural *Bacillus* spp. isolates in order to find prospective strains which could be possibly used in QQ-mediated control of AHL-producing phytopathogens. The collection of 149 *Bacillus* spp. isolates, originated from soil, manure and straw, was screened for the presence and activity of AHL-lactonases using well-diffusion method. Violacein production by biosensor strain *Chromobacterium violaceum* CV026 was indicator for AHL-presence and degradation. PCR method was performed for the detection of *aiiA* gene in selected isolates, with primers *aiiA1-F* and *aiiA2-R*. Degradation of AHLs in medium was observed by the loss of violacein purple colour around wells. Total of 52 strains showed AHL-lactonase activity, which was confirmed by detection of a single 793 bp long band of amplified *aiiA* gene. Furthermore, difference in AHL-lactonase production was perceived in *Bacillus* spp. isolates from different natural samples. The highest number of AHL-lactonase producing *Bacillus* spp. were found in soil samples (65 %), followed by manure samples (33 %), while only one strain from straw demonstrated AHL-lactonase activity. The results suggest that AHL-lactonase is common in *Bacillus* spp. exceptionally in soil isolates.

Keywords: *Bacillus* spp.; biocontrol; *quorum quenching*; AHL-lactonase activity.

V. major onto leaves of five healthy plants. Five non-inoculated plants served as controls. Inoculated and non-inoculated plants were kept in a greenhouse at 24 to 30°C. Inoculated plants developed signs and symptoms after 10 days, control plants remained symptomless. The fungus from the inoculated plants was identical morphologically to that observed from initially diseased plants. To confirm the identification, the internal transcribed spacer (ITS) regions of PMV12 were amplified with primers ITS1/4 and ITS5/P3 and sequenced directly. The resulting sequences were deposited in GenBank (Accession Nos. OQ165183 for ITS1/4 and OQ165185 for ITS5/P3). BLAST search of these sequences revealed 100% similarity with the ITS sequences of *Golovinomyces orontii* on plants of *Vinca* sp. (KY660780) and *Vinca major* (KR011138). Based on the morphological characteristics and molecular analysis, the fungus was identified as *G. orontii*. To our knowledge, this is the first report of powdery mildew caused by *G. orontii* on *V. major* in Türkiye.

Key words: *Golovinomyces orontii*, *Vinca major*, powdery mildew

P-9-STU

Population dynamics of *Bacillus amyloliquefaciens* SS-38.4 in the phyllosphere of sugar beet and its biocontrol activity against *Pseudomonas syringae* pv. *aptata* P21

Iva Rosić*, Marina Anteljević, Tamara Ranković, Ivan Nikolić, Slaviša Stanković, Tanja Berić, Olja Medić

University of Belgrade, Faculty of Biology, Center for Biological Control and Plant Growth Promotion, 11000 Belgrade, Serbia

* Correspondence: iva.rosic@bio.bg.ac.rs

The success of biocontrol depends directly on the colonization of plant tissue by the biocontrol agent. The objective of this study was to investigate the ability of *Bacillus amyloliquefaciens* SS-38.4 to colonize sugar beet leaves and suppress the leaf spot disease caused by *Pseudomonas syringae* pv. *aptata* P21. Sugar beet leaves were sprayed with the bacterial suspension of SS-38.4 followed by P21 12 hours later (T_1), but also in the reverse order (T_2). Plants treated with SS-38.4 or P21 alone were controls ($C_{38.4}$ and C_{P21}). Strains were isolated one, five and seven days after treatments and confirmed with SS-38.4-SCAR markers and *P. syringae*-specific primers. Results were analysed using two-way ANOVA and Tukey's test ($\alpha = 0.05$). During seven days, populations were stable in both controls ($C_{38.4}$ 3.45 ± 0.31 and C_{P21} 3.72 ± 0.44 log CFU/cm²). Changes in population number of P21 were noticeable from the fifth day (T_1 1.35 and T_2 1.43 log CFU/cm²), while on the seventh day a significant decrease was observed in T_1 (0 log CFU/cm²) and T_2 (1 log CFU/cm²). The number of SS-38.4 remained constant in both treatments and similar to control. A significant reduction in disease symptoms was observed on the seventh day in T_1 and T_2 (0 % and 0.23% affected leaf tissue) compared to C_{P21} (3.9 %). Strain *B. amyloliquefaciens* SS-38.4 can establish and maintain a stable population on the sugar beet leaf surface up to seven days after application, while exhibiting a significant suppressive effect on *P. syringae* pv. *aptata* P21.

Key words: Biocontrol, *Bacillus* spp., *Pseudomonas* spp., phytopathogen, phyllosphere colonization

P-10-STU

Phylogenetic analysis of *Pseudomonas syringae* isolates from the Danube River Basin revealed association with past epidemics in Serbia

Marina Anteljević*, Iva Rosić, Tamara Ranković, Olja Medić, Tanja Berić, Slaviša Stanković, Ivan Nikolić

University of Belgrade, Faculty of Biology, Center for Biological Control and Plant Growth Promotion, 11000 Belgrade, Serbia

*Correspondence: marina.anteljevic@bio.bg.ac.rs

Pseudomonas syringae (*Psy*) is a widespread complex of plant pathogenic bacteria. It's a causal agent of diseases in many economically important hosts, including herbaceous and woody plants. The species complex is composed of 13 phylogroups. The aim of our study was to estimate the phylogenetic diversity of *Psy* isolates from the Serbian Danube River basin – a major irrigation source. A partial sequence of the citrate synthase housekeeping gene (*cts*) was amplified for 51 isolates from the collection. All amplicons were sequenced using the Sanger sequencing method. Based on the sequences obtained, a phylogenetic tree was constructed using Mega 11 software to infer the evolutionary history using the neighbour-joining method. The *cts* sequences of 51 isolates were compared with the *cts* sequences of isolates from previously reported epidemics in Serbia using the NCBI BLASTn tool. The analysis resulted in the detection of phylogroups 2, 7, 9, and 13. Most isolates were assigned to phylogroup 2 (70.6%), with the remainder evenly distributed among phylogroups 7, 9, and 13 (9.8% each). Nine isolates of phylogroup 2 showed 100% similarity in *cts* sequence with isolates from diseased cherry plants in different epidemic events in Serbia. Phylogroup 2 is known to be the most widespread phylogroup, of which numerous isolates have been found in non-agricultural habitats but also as disease causal agents on plants. Insights into phylogenetic diversity in the environment are important to explain the ecology of *Psy* and to predict possible disease outbreaks.

Key words: *Pseudomonas syringae*, phylogeny, diversity, species complex, citrate-synthase gene

Заступљеност биљног патогена *Pseudomonas syringae* у Увачким језерима

Марина Антељевић¹, Иван Николић¹, Стоимир Коларевић², Ива Росић¹, Тамара Павловић¹, Тања Берић¹, Славиша Станковић¹

¹Универзитет у Београду – Биолошки факултет, Катедра за микробиологију, Београд, Србија
marina.anteljevic@bio.bg.ac.rs

²Институт за биолошка истраживања „Синиша Станковић“ – Институт од националног значаја за Републику Србију, Универзитет у Београду, Београд, Србија

Pseudomonas syringae је један од најбоље проучених биљних патогена који инфицира готово све економски важне културе.¹ Овај патоген опстаје и размножава се у различитим стаништима повезаним са циклусом кружења воде у природи.² У циљу утврђивања заступљености *P. syringae* у Увачким језерима узорковање је вршено крајем августа 2021. године на четири локалитета на различитим дубинама: Ушће Кладнице (0,5 m и 5 m), Радоињско језеро (0,5 m, 9 m и 12 m), Сјеничко језеро (0,5 m и 5 m) и Златарско језеро (0,5 m, 5 m и 8 m). Урађена је мембранска филтрација узорака воде, а филтери су пребачени на хранљиву подлогу селективну за род *Pseudomonas*. Коришћењем прајмера дизајнираних за детекцију *P. syringae*³ изолати су идентификовани у *colony*-PCR реакцији. Заступљеност *P. syringae* у Радоињском језеру на дубини 9 m била је 10 cfu/L, а на дубини 12 m 15 cfu/L. У Сјеничком језеру на дубини 5 m заступљеност је била 5 cfu/L. У Златарском језеру на дубини 0,5 m било је присутно 5 cfu/L, док је на дубини 8 m било 25 cfu/L. У односу на укупан број израслих *Pseudomonas* колонија проценат заступљености *P. syringae* се кретао између 2,25% и 5,56%. Без обзира на ниску заступљеност, карактеризација изолата и детектовање потенцијалног извора инфекције изван агроекосистема је битна мера у превенцији нових инфекција и спречавању смањења пољопривредних приноса.

1. Xin, X.F., Kvitko, B., He, S.Y., 2018, Nat. Rev. Microbiol. 16:316-328.
2. Morris, C.E., Monteil, C.L., Berge, O., 2013, Annu. Rev. Phytopathol. 51:85-104.
3. Guilbaud, C., Morris, C.E., Barakat, M., et al., 2016, FEMS Microbiol. Ecol. 92:fiv146.

Захвалница: Овај рад је финансиран од стране Министарства просвете, науке и технолошког развоја Републике Србије, Уговори бр. 451-03-68/2022-14/200178 и 451-03-9/2021-14/200007.

In vitro* испитивање потенцијала липопептидних екстраката изолата *Bacillus* spp. за сузбијање биљног патогена *Pseudomonas syringae

Ива Росић, Иван Николић, Оља Медић, Тамара Павловић, Марина Антељевић, Тања Берић, Славиша Станковић

Универзитет у Београду – Биолошки факултет, Катедра за микробиологију, Београд, Србија, iva.rosic@bio.bg.ac.rs

Бактерије рода *Bacillus* поседују велики биоконтролни потенцијал захваљујући продукцији липопептида са израженом активношћу против различитих фитопатогена. Примена липопептида, као алтернативе хемијским пестицидима, омогућава превазилажење проблема резистенције биљних патогена и загађења животне средине.¹ *Pseudomonas syringae* представља економски важног патогена услед неповољног утицаја на велики број различитих пољопривредних култура.² Циљ овог истраживања је био *in vitro* испитивање потенцијала 246 изолата рода *Bacillus* у инхибицији *P. syringae* P17 и P21 сојева. Применом *well-diffusion* методе са преконоћним културама и супернатантима *Bacillus* spp. изолата, одабрано је шест изолата са највећим пречницима зона инхибиције раста (≥ 17 mm) оба испитивана *P. syringae* соја. Након етил-ацетатне екстракције липопептида испитане су минималне инхибиторне концентрације (МИК) и минималне бактерицидне концентрације (МБК) шест екстраката. Најефикасније се показао сој SS-8.1.2, са МИК 0,3125 mg/ml и МБК 5 mg/ml. Потенцијални механизам деловања антимикуробних једињења, синергизам,³ испитан је мешањем екстракта SS-8.1.2 са екстрактом другог изолата по ефикасности (SS-36.3 са МИК 5 mg/ml) у односу 1:1. Комбинација је показала антагонистичку активност на основу индекса фракционе инхибиторне концентрације (ФИЦИ ≥ 2). Добијени резултати показали су инхибиторно дејство липопептидних екстраката појединих *Bacillus* spp. изолата и отворили могућност за даље тестове *in planta* применом SS-8.1.2 екстракта.

1. Penha, R.O., Vandenberghe, L.P.S., Faulds, C., *et al.*, 2020, *Planta* 251:70.
2. Nikolić, I., Berić, T., Dimkić, I., *et al.*, 2019, *J. Appl. Microbiol.* 126:165-176.
3. Dimkić, I., Stanković, S., Nišavić, M., *et al.*, 2017, *Front. Microbiol.* 8:925.

Захвалница: Овај рад је финансиран од стране Министарства просвете, науке и технолошког развоја Републике Србије, Уговор бр. 451-03-68/2022-14/200178.

S5A-P5
Phylogenomic status of two *P. syringae* strains P16 and P21 with different pathogenicity isolated from sugar beet in Serbia

Nikolić I.¹, Pavlović T.¹, Rosić I.¹, Anteljević M.¹, Medić O.¹, Berić T.¹ & Stanković S.¹

¹ Faculty of Biology, Center for biocontrol and plant growth promotion, University of Belgrade, Belgrade, Serbia
 E-mail: ivan.nikolic@bio.bg.ac.rs

Members of the *Pseudomonas syringae* species complex are heterogeneous bacteria with strong abilities to exist on and infect different plant hosts and survive beyond agroecosystems^[1]. The *P. syringae* strains previously collected from sugar beet commercial fields in Serbia were characterized as a genetically and pathogenically diverse strains collection^[2]. This study provides comparative genomics and phylogenomic status assessment of two *P. syringae* strains from this collection, P16 and P21, with distinctive pathogenicity. The whole-genome data obtained in this study were combined with the publicly available genomes of 34 strains from the *P. syringae* species complex, including strains from the most ubiquitous and pathogenic phylogroups and strains isolated from non-agricultural environments. The core SNP and ANI analysis revealed four distinctive clusters among compared genomes, where P16 and P21 were clustered on the same branch, mostly with strains from PG02 and two strains from PG01 (*P. syringae* isolated from kiwi fruit). The pangenome analysis revealed 6.03% of the core genome and 93.97% of accessory genes. Based on the presence/absence of individual genes, both strains are grouped in the same branch of the phylogenomic tree together with other *P. syringae* strains from PG02b. We observed slightly different virulence factor gene composition between P16 and P21 strains, especially the T3SS effector and T4SS *virB* genes repertoire. The extensive accessory genome revealed a high degree of variability among *P. syringae* complex, which could be associated with their ability to survive in diverse ecological niches and the extensive horizontal gene transfer^[3].

^[1]Xin *et al.*, 2018. Nat. Rev. Microbiol., 16, 316–328

^[2]Nikolić *et al.*, 2018 Plant Pathol., 67(5), 1194-1207

^[3]Dillon *et al.*, 2019. Genome Biol., 20: 1-28.

S5A-P6

Killing effect of *Bacillus velezensis* FZB42 on a *Xanthomonas campestris* pv. *campestris* strain newly isolated from cabbage: a metabolomic study

Novotny C.^{1,3}, Maresova H.², Macha H.², Benada O.², Palyzova A.²

¹ Laboratory of Environmental Biotechnology, Institute of Microbiology of the CAS, Videnska 1083, 142 20 Prague 4, Czech Republic

² Laboratory of Characterization of Molecular Structures, Institute of Microbiology of the CAS, Videnska 1083, 142 20 Prague 4, Czech Republic

³ Department of Horticulture, Faculty of Agrobiology, Food and Natural Resources, Czech University of Life Sciences, Prague, Kamýcka 129, 165 21 Prague 6, Czech Republic

E-mail: novotny@biomed.cas.cz

Potential of *Bacillus velezensis* for biological control of various phytopathogens has been documented over the past few years but its antagonistic interactions with xanthomonads have not been studied in detail. The findings documented a strong killing effect on *Xanthomonas campestris* pv. *campestris* (Xcc) cells in a co-culture with *B. velezensis*. Lipopeptides and the siderophore bacillibactin involved in the killing process were quantified. A new robust Xcc-SU isolate tolerating high concentrations of ferric ions was used. In a co-culture with the antagonist, the population of Xcc-SU was annihilated within 24–48 h, depending on the number of antagonist cells used for inoculation. No inhibitory effect of Xcc-SU on *B. velezensis* was observed. Bacillibactin and lipopeptides (surfactin, fengycin, bacillomycin) were present in both the co-culture and the monoculture of *B. velezensis*. Except for bacillibactin, the maximum contents of lipopeptides were higher in the antagonist monoculture compared with the co-culture. Scanning electron microscopy showed that the death of Xcc-SU bacteria in the co-culture was caused by cell lysis. The analysis by mass spectrometry showed four major compounds, bacillibactin, surfactin, fengycin, and bacillomycin D. Different forms of surfactin and fengycin with variations in their side-chain length were detected. The results demonstrated the ability of *B. velezensis* FZB42 to act as a powerful antagonistic strain against Xcc.

Funding information: The work was supported by the following projects: EU COST No. CA16107, INTER-COST LTC18009, RVO 61388971 (Ministry of Education, Youth and Sports of the Czech Republic), 19-10907S (Czech Science Foundation), IGA PrF 2021 021 (Palacký University, Czech Republic), QK1910235 (Ministry of Agriculture of the Czech Republic).



УНИВЕРЗИТЕТ У БЕОГРАДУ
БИОЛОШКИ ФАКУЛТЕТ

Студентски трг 16
11000 БЕОГРАД
Република СРБИЈА
Тел: +381 11 2186 635
Факс: +381 11 2638 500
Е-пошта: dekanat@bio.bg.ac.rs

Број: пБ3003/2021-8
Београд, 28. децембар 2023.
Број индекса: Б3003/2021

Универзитет у Београду, Биолошки факултет, на захтев који је поднела студенткиња **Ива Росић** издаје следећу

ПОТВРДУ

Студенткиња **Ива Росић**, број индекса **Б3003/2021**, рођена **9. маја 1997**, уписала је школску **2023/2024.** годину на Универзитету у Београду, Биолошком факултету, студијски програм **Биологија**, модул **Биологија микроорганизама - докторске академске студије**, као студент који се финансира из буџета Републике Србије.

Студенткиња је уписана први пут на Универзитет у Београду, Биолошки факултет, студијски програм **Биологија**, модул **Биологија микроорганизама - докторске академске студије** школске **2021/2022.** године.

Након завршетка докторских академских студија студенткиња ће стећи научни назив доктор наука - биолошке науке.

Студенту је на II редовној седници Наставно-научног већа Биолошког факултета Универзитета у Београду одржаној 10. 11. 2023. године прихваћена пријава теме за израду докторске дисертације.

Потврда се издаје на лични захтев ради **регулisaња избора у звање** и не може се користити у друге сврхе.

ОВЛАШЋЕНО ЛИЦЕ СТУДЕНТСКЕ СЛУЖБЕ БИОЛОШКОГ ФАКУЛТЕТА



Зоран Николић



Република Србија
Универзитет у Београду

Оснивач: Република Србија

Дозволу за рад број 612-00-02666/2010-04 од 12. октобра 2011.
године је издало Министарство просвете и науке Републике Србије

Биолошки факултет, Београд

Оснивач: Република Србија

Дозволу за рад број 612-00-01196/2009-04 од 24. децембра 2010.
године је издало Министарство просвете Републике Србије

УБ



Диплома

Ива, Мирко, Росић

рођена 9. маја 1997. године, Ужице, Република Србија, уписана школске
2016/2017. године, а дана 13. јула 2020. године завршила је основне академске
студије, првог степена, на студијском програму Биологија, обима 240
(двеста четрдесет) бодова ЕСПБ са просечном оценом 9,60 (девет и 60/100).

На основу тога издаје јој се ова диплома о стеченом високом образовању и стручном називу
дипломирани биолог

Број: 11017000

У Београду, 30. септембра 2020. године

Декан
Проф. др Желко Томановић

Ректор
Проф. др Иванка Појовић

00110341



Република Србија
Универзитет у Београду

Оснивач: Република Србија
Дозвола за рад број 612-00-02666/2010-04 од 12. октобра 2011.
године је издало Министарство просвете и науке Републике Србије

Биолошки факултет, Београд

Оснивач: Република Србија
Дозвола за рад број 612-00-01196/2009-04 од 24. децембра 2010.
године је издало Министарство просвете Републике Србије

УБ

Диплома

Ива, Мирко, Росић

рођена 9. маја 1997. године, Ужице, Република Србија, уписана школске 2020/2021.

године, а дана 16. јула 2021. године завршила је мастер академске студије,
групе степен, на студијском програму Молекуларна биологија и физиологија,
обима 60 (шездесет) бодова ЕСПБ са просечном оценом 10,00 (десет и 0/100).

На основу тога издаје јој се ова диплома о стеченом високом образовању и академском називу

мастер биологије

Број: 12570400

У Београду, 6. октобра 2021. године

Декан
Проф. др Љубиша Станисављевић

[Signature]

Ректор
Проф. др Владан Ђокић

[Signature]

00125878



УНИВЕРЗИТЕТ У БЕОГРАДУ

Адреса: Студентски трг 1, 11000 Београд, Република Србија
Тел.: 011 3207400; Факс: 011 2638818; E-mail: kabinet@rect.bg.ac.rs

ВЕЋЕ НАУЧНИХ ОБЛАСТИ
ПРИРОДНИХ НАУКА

Београд, 21. децембар 2023. године
02-07 Број: 61206-4683/2-23
МЦ

На основу члана 48 став 5 тачка 3 Статута Универзитета у Београду („Гласник Универзитета у Београду“, бр. 201/18, 207/19, 213/20, 214/20, 217/20, 230/21, 232/22, 233/22, 236/22, 241/22, 243/22, 244/23 245/23 и 247/23) и члана 32 Правилника о докторским студијама на Универзитету у Београду („Гласник Универзитета у Београду“, бр. 191/2016, 212/2019, 215/2020, 217/2020, 228/21, 230/21 и 241/22), а на захтев Биолошког факултета, бр. 50/214-1 од 12. децембра 2023. године, Веће научних области природних наука, на седници одржаној 21. децембра 2023. године, донело је

О Д Л У К У

ДАЈЕ СЕ САГЛАСНОСТ на одлуку Наставно-научног већа Биолошког факултета о прихватању теме докторске дисертације ИВЕ РОСИЋ, под називом: „Фенотипска и генотипска анализа капацитета *Vacillus* spp. изолата за колонизацију, поспешивање раста биљака и биоконтролу“ и одређивању др Оље Медић, доцента Универзитета у Београду – Биолошког факултета за ментора.

ПРЕДСЕДНИЦА ВЕЋА

проф. др Јасмина Димитрић-Марковић

Доставити:

- Факултету
- архиви Универзитета